

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number
WO 01/21810 A1

(51) International Patent Classification⁷: C12N 15/54,
15/62, 15/85, 9/10, C07K 16/40, A61K 31/711, 38/45,
39/40, G01N 33/53

(74) Agents: NGUYEN, Thuy, H. et al.; Smart & Biggar,
900-55 Metcalfe Street, P.O. Box 2999, Station D, Ottawa,
Ontario K1P 5Y6 (CA).

(21) International Application Number: PCT/CA00/01085

(22) International Filing Date:
15 September 2000 (15.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/154,461 17 September 1999 (17.09.1999) US

(71) Applicant (for all designated States except US): AVEN-
TIS PASTEUR LIMITED [CA/CA]; Connaught Cam-
pus, 1755 Steeles Avenue West, Toronto, Ontario M2R 3T4
(CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MURDIN, Andrew,
D. [CA/CA]; 11 Forest Hill Drive, Richmond Hill L4B
3C2 (CA). OOMEN, Raymond, P. [CA/CA]; 29 Kennedy
St. W., Aurora, Ontario L4G 2L6 (CA). WANG, Joe
[CA/CA]; 48 29th Street, Etobicoke, Ontario M8W 3A8
(CA). DUNN, Pamela [GB/CA]; Apartment 703, 370
Kaneff Crescent, Mississauga, Ontario L5A 4B8 (CA).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: *CHLAMYDIA* ANTIGENS AND CORRESPONDING DNA FRAGMENTS AND USES THEREOF

(57) Abstract: The present invention provides a method of nucleic acid, including DNA, immunization of a host, including humans, against disease caused by infection by a strain of *Chlamydia*, specifically *C. pneumoniae*, employing a vector containing a nucleotide sequence encoding lpxB of a strain of *Chlamydia pneumoniae* and a promoter to effect expression of the lpxB gene in the host. Modifications are possible within the scope of this invention.



WO 01/21810 A1

TITLE OF INVENTION

CHLAMYDIA ANTIGENS AND CORRESPONDING DNA
FRAGMENTS AND USES THEREOF

REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of U.S.
Provisional Application No. 60/154,461, filed September 17,
1999.

FIELD OF INVENTION

10 The present invention relates to the *Chlamydia* lpxB
protein antigen and corresponding DNA molecules, which can be
used to prevent and treat *Chlamydia* infection in mammals, such
as humans.

BACKGROUND OF THE INVENTION

15 Chlamydiae are prokaryotes. They exhibit morphologic
and structural similarities to gram-negative bacteria including
a trilaminar outer membrane, which contains lipopolysaccharide
and several membrane proteins that are structurally and
functionally analogous to proteins found in *E coli*. They are
obligate intra-cellular parasites with a unique biphasic life
20 cycle consisting of a metabolically inactive but infectious
extracellular stage and a replicating but non-infectious
intracellular stage. The replicative stage of the life-cycle
takes place within a membrane-bound inclusion which sequesters
the bacteria away from the cytoplasm of the infected host cell.

25 *C. pneumoniae* is a common human pathogen, originally
described as the TWAR strain of *Chlamydia psittaci* but
subsequently recognised to be a new species. *C. pneumoniae* is
antigenically, genetically and morphologically distinct from
other chlamydia species (*C. trachomatis*, *C. pecorum* and *C.*
30 *psittaci*). It shows 10% or less DNA sequence homology with
either of *C. trachomatis* or *C. psittaci*.

C. pneumoniae is a common cause of community acquired pneumonia, only less frequent than *Streptococcus pneumoniae* and *Mycoplasma pneumoniae* (Grayston et al. (1995) Journal of Infectious Diseases 168:1231; Campos et al. (1995) Investigation of Ophthalmology and Visual Science 36:1477). It can also cause upper respiratory tract symptoms and disease, including bronchitis and sinusitis (Grayston et al. (1995) Journal of Infectious Diseases 168:1231; Grayston et al (1990) Journal of Infectious Diseases 161:618; Marrie (1993) Clinical Infectious Diseases. 18:501; Wang et al (1986) Chlamydial infections Cambridge University Press, Cambridge. p. 329. The great majority of the adult population (over 60%) has antibodies to *C. pneumoniae* (Wang et al (1986) Chlamydial infections. Cambridge University Press, Cambridge. p. 329), indicating past infection which was unrecognized or asymptomatic.

C. pneumoniae infection usually presents as an acute respiratory disease (i.e., cough, sore throat, hoarseness, and fever; abnormal chest sounds on auscultation). For most patients, the cough persists for 2 to 6 weeks, and recovery is slow. In approximately 10% of these cases, upper respiratory tract infection is followed by bronchitis or pneumonia. Furthermore, during a *C. pneumoniae* epidemic, subsequent co-infection with pneumococcus has been noted in about half of these pneumonia patients, particularly in the infirm and the elderly. As noted above, there is more and more evidence that *C. pneumoniae* infection is also linked to diseases other than respiratory infections.

The reservoir for the organism is presumably people. In contrast to *C. psittaci* infections, there is no known bird or animal reservoir. Transmission has not been clearly defined. It may result from direct contact with secretions, from fomites, or from airborne spread. There is a long incubation period, which may last for many months. Based on

analysis of epidemics, *C. pneumoniae* appears to spread slowly through a population (case-to-case interval averaging 30 days) because infected persons are inefficient transmitters of the organism. Susceptibility to *C. pneumoniae* is universal.

5 Reinfections occur during adulthood, following the primary infection as a child. *C. pneumoniae* appears to be an endemic disease throughout the world, noteworthy for superimposed intervals of increased incidence (epidemics) that persist for 2 to 3 years. *C. trachomatis* infection does not confer cross-
10 immunity to *C. pneumoniae*. Infections are easily treated with oral antibiotics, tetracycline or erythromycin (2 g/d, for at least 10 to 14 d). A recently developed drug, azithromycin, is highly effective as a single-dose therapy against chlamydial infections.

15 In most instances, *C. pneumoniae* infection is often mild and without complications, and up to 90% of infections are subacute or unrecognized. Among children in industrialized countries, infections have been thought to be rare up to the age of 5 y, although a recent study (E Normann et al, Chlamydia
20 pneumoniae in children with acute respiratory tract infections, Acta Paediatrica, 1998, Vol 87, Iss 1, pp 23-27) has reported that many children in this age group show PCR evidence of infection despite being seronegative, and estimates a prevalence of 17-19% in 2-4 y olds. In developing countries,
25 the seroprevalence of *C. pneumoniae* antibodies among young children is elevated, and there are suspicions that *C. pneumoniae* may be an important cause of acute lower respiratory tract disease and mortality for infants and children in tropical regions of the world.

30 From seroprevalence studies and studies of local epidemics, the initial *C. pneumoniae* infection usually happens between the ages of 5 and 20 y. In the USA, for example, there are estimated to be 30,000 cases of childhood pneumonia each year caused by *C. pneumoniae*. Infections may cluster among

groups of children or young adults (e.g., school pupils or military conscripts).

C. pneumoniae causes 10 to 25% of community-acquired lower respiratory tract infections (as reported from Sweden, Italy, Finland, and the USA). During an epidemic, *C. pneumonia* infection may account for 50 to 60% of the cases of pneumonia. During these periods, also, more episodes of mixed infections with *S. pneumoniae* have been reported.

Reinfection during adulthood is common; the clinical presentation tends to be milder. Based on population seroprevalence studies, there tends to be increased exposure with age, which is particularly evident among men. Some investigators have speculated that a persistent, asymptomatic *C. pneumoniae* infection state is common.

In adults of middle age or older, *C. pneumoniae* infection may progress to chronic bronchitis and sinusitis. A study in the USA revealed that the incidence of pneumonia caused by *C. pneumoniae* in persons younger than 60 years is 1 case per 1,000 persons per year; but in the elderly, the disease incidence rose three-fold. *C. pneumoniae* infection rarely leads to hospitalization, except in patients with an underlying illness.

Of considerable importance is the association of atherosclerosis and *C. pneumoniae* infection. There are several epidemiological studies showing a correlation of previous infections with *C. pneumoniae* and heart attacks, coronary artery and carotid artery disease (Saikku et al. (1988) Lancet;ii:983; Thom et al. (1992) JAMA 268:68; Linnanmaki et al. (1993), Circulation 87:1030; Saikku et al. (1992) Annals Internal Medicine 116:273; Melnick et al (1993) American Journal of Medicine 95:499). Moreover, the organisms has been detected in atheromas and fatty streaks of the coronary, carotid, peripheral arteries and aorta (Shor et al. (1992) South African. Medical Journal 82:158; Kuo et al. (1993) Journal of

Infectious Diseases 167:841; Kuo et al. (1993) Arteriosclerosis and Thrombosis 13:1500; Campbell et al (1995) Journal of Infectious Diseases 172:585; Chiu et al. Circulation, 1997 (In Press)). Viable *C. pneumoniae* has been recovered from the coronary and carotid artery (Ramirez et al (1996) Annals of Internal Medicine 125:979; Jackson et al. Abst. K121, p272, 36th ICAAC, 15-18 Sept. 1996, New Orleans). Furthermore, it has been shown that *C. pneumoniae* can induce changes of atherosclerosis in a rabbit model (Fong et al (1997) Journal of Clinical Microbiology 35:48). Taken together, these results indicate that it is highly probable that *C. pneumoniae* can cause atherosclerosis in humans, though the epidemiological importance of chlamydial atherosclerosis remains to be demonstrated.

15 A number of recent studies have also indicated an association between *C. pneumoniae* infection and asthma. Infection has been linked to wheezing, asthmatic bronchitis, adult-onset asthma and acute exacerbations of asthma in adults, and small-scale studies have shown that prolonged antibiotic treatment was effective at greatly reducing the severity of the disease in some individuals (Hahn DL, et al. Evidence for Chlamydia pneumoniae infection in steroid-dependent asthma. Ann Allergy Asthma Immunol. 1998 Jan; 80(1): 45-49.; Hahn DL, et al. Association of Chlamydia pneumoniae IgA antibodies with recently symptomatic asthma. Epidemiol Infect. 1996 Dec; 20 117(3): 513-517; Bjornsson E, et al. Serology of chlamydia in relation to asthma and bronchial hyperresponsiveness. Scand J Infect Dis. 1996; 28(1): 63-69.; Hahn DL. Treatment of Chlamydia pneumoniae infection in adult asthma: a before-after trial. J Fam Pract. 1995 Oct; 41(4): 345-351.; Allegra L, et al. Acute exacerbations of asthma in adults: role of Chlamydia pneumoniae infection. Eur Respir J. 1994 Dec; 7(12): 2165-2168.; Hahn DL, et al. Association of Chlamydia pneumoniae

(strain TWAR) infection with wheezing, asthmatic bronchitis, and adult-onset asthma. JAMA. 1991 Jul 10; 266(2): 225-230).

In light of these results a protective vaccine against *C. pneumoniae* infection would be of considerable importance. There is not yet an effective vaccine for any human chlamydial infection. It is conceivable that an effective vaccine can be developed using physically or chemically inactivated Chlamydiae. However, such a vaccine does not have a high margin of safety. In general, safer vaccines are made by genetically manipulating the organism by attenuation or by recombinant means. Accordingly, a major obstacle in creating an effective and safe vaccine against human chlamydial infection has been the paucity of genetic information regarding Chlamydia, specifically *C. pneumoniae*.

Studies with *C. trachomatis* and *C. psittaci* indicate that safe and effective vaccine against Chlamydia is an attainable goal. For example, mice which have recovered from a lung infection with *C. trachomatis* are protected from infertility induced by a subsequent vaginal challenge (Pal et al. (1996) Infection and Immunity.64:5341). Similarly, sheep immunized with inactivated *C. psittaci* were protected from subsequent chlamydial-induced abortions and stillbirths (Jones et al. (1995) Vaccine 13:715). Protection from chlamydial infections has been associated with Th1 immune responses, particularly the induction of INF γ - producing CD4+T-cells (Igiertsemes et al. (1993) Immunology 5:317). The adoptive transfer of CD4+ cell lines or clones to nude or SCID mice conferred protection from challenge or cleared chronic disease (Igiertseme et al (1993) Regional Immunology 5:317; Magee et al (1993) Regional Immunology 5: 305), and *in vivo* depletion of CD4+ T cells exacerbated disease post-challenge (Landers et al (1991) Infection & Immunity 59:3774; Magee et al (1995) Infection & Immunity 63:516). However, the presence of sufficiently high titres of neutralising antibody at mucosal

surfaces can also exert a protective effect (Cotter et al. (1995) Infection and Immunity 63:4704).

Antigenic variation within the species *C. pneumoniae* is not well documented due to insufficient genetic information, though variation is expected to exist based on *C. trachomatis*. Serovars of *C. trachomatis* are defined on the basis of antigenic variation in the major outer membrane protein (MOMP), but published *C. pneumoniae* MOMP gene sequences show no variation between several diverse isolates of the organism (Campbell et al. Infection and Immunity (1990) 58:93; McCafferty et al Infection and Immunity (1995) 63:2387-9; Gaydos et al. Infection and Immunity.(1992) 60(12):5319-5323). The gene encoding a 76 kDa antigen has been cloned from a single strain of *C. pneumoniae* and the sequence published (Perez Melgosa et al. Infection and Immunity.(1994) 62:880). An operon encoding the 9 kDa and 60 kDa cyteine-rich outer membrane protein genes has been described (Watson et al., Nucleic Acids Res (1990) 18:5299; Watson et al., Microbiology (1995) 141:2489). Many antigens recognized by immune sera to *C. pneumoniae* are conserved across all chlamydiae, but 98 kDa, 76 kDa and several other proteins may be *C. pneumoniae*-specific (Perez Melgosa et al. Infection and Immunity.(1994) 62:880; Melgosa et al., FEMS Microbiol Lett (1993) 112 :199;, Campbell et al., J Clin Microbiol (1990) 28 :1261; Iijima et al., J Clin Microbiol (1994) 32:583). An assessment of the number and relative frequency of any *C. pneumoniae* serotypes, and the defining antigens, is not yet possible. The entire genome sequence of *C. pneumoniae* strain CWL-029 is now known (<http://chlamydia-www.berkeley.edu:4231/>) and as further sequences become available a better understanding of antigenic variation may be gained.

Many antigens recognised by immune sera to *C. pneumoniae* are conserved across all chlamydiae, but 98kDa, 76 kDa and 54 kDa proteins appear to be *C. pneumoniae*-specific

(Campos et al. (1995) Investigation of Ophthalmology and Visual Science 36:1477; Marrie (1993) Clinical Infectious Diseases. 18:501; Wiedmann-Al-Ahmad M, et al. Reactions of polyclonal and neutralizing anti-p54 monoclonal antibodies with an isolated, species-specific 54-kilodalton protein of *Chlamydia pneumoniae*. Clin Diagn Lab Immunol. 1997 Nov; 4(6): 700-704).

Immunoblotting of isolates with sera from patients does show variation of blotting patterns between isolates, indicating that serotypes *C. pneumoniae* may exist (Grayston et al. (1995) Journal of Infectious Diseases 168:1231; Ramirez et al (1996) Annals of Internal Medicine 125:979). However, the results are potentially confounded by the infection status of the patients, since immunoblot profiles of a patient's sera change with time post-infection. An assessment of the number and relative frequency of any serotypes, and the defining antigens, is not yet possible.

Accordingly, a need exists for identifying and isolating polynucleotide sequences of *C. pneumoniae* for use in preventing and treating *Chlamydia* infection.

SUMMARY OF THE INVENTION

The present invention provides purified and isolated polynucleotide molecules that encode the *Chlamydia* polypeptides designated lpxB (SEQ ID No: 1) which can be used in methods to prevent, treat, and diagnose *Chlamydia* infection. In one form of the invention, the polynucleotide molecules are DNA that encode the polypeptide of SEQ ID No: 2.

Another form of the invention provides polypeptides corresponding to the isolated DNA molecules. The amino acid sequence of the corresponding encoded polypeptide is shown as SEQ ID No: 2.

Those skilled in the art will readily understand that the invention, having provided the polynucleotide sequences encoding the *Chlamydia* lpxB protein, also provides

polynucleotides encoding fragments derived from such a polypeptide. Moreover, the invention is understood to provide mutants and derivatives of such polypeptides and fragments derived therefrom, which result from the addition, deletion, or substitution of non-essential amino acids as described herein. Those skilled in the art would also readily understand that the invention, having provided the polynucleotide sequences encoding *Chlamydia* polypeptides, further provides monospecific antibodies that specifically bind to such polypeptides.

The present invention has wide application and includes expression cassettes, vectors, and cells transformed or transfected with the polynucleotides of the invention. Accordingly, the present invention further provides (i) a method for producing a polypeptide of the invention in a recombinant host system and related expression cassettes, vectors, and transformed or transfected cells; (ii) a vaccine, or a live vaccine vector such as a pox virus, *Salmonella typhimurium*, or *Vibrio cholerae* vector, containing a polynucleotide of the invention, such vaccines and vaccine vectors being useful for, e.g., preventing and treating *Chlamydia* infection, in combination with a diluent or carrier, and related pharmaceutical compositions and associated therapeutic and/or prophylactic methods; (iii) a therapeutic and/or prophylactic use of an RNA or DNA molecule of the invention, either in a naked form or formulated with a delivery vehicle, a polypeptide or combination of polypeptides, or a monospecific antibody of the invention, and related pharmaceutical compositions; (iv) a method for diagnosing the presence of *Chlamydia* in a biological sample, which can involve the use of a DNA or RNA molecule, a monospecific antibody, or a polypeptide of the invention; and (v) a method for purifying a polypeptide of the invention by antibody-based affinity chromatography.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention will be further understood from the following description with reference to the drawings, in which:

5 Figure 1 shows the nucleotide sequence of the lpxB gene (SEQ ID No: 1) and the deduced amino acid sequence of lpxB from *Chlamydia pneumoniae* (SEQ ID No: 2).

 Figure 2 shows the restriction enzyme analysis of the *C. pneumoniae* lpxB gene.

10 Figure 3 shows the construction and elements of plasmid pCABk1043.

 Figure 4 illustrates protection against *C. pneumoniae* infection by pCABk1043 following DNA immunization.

DETAILED DESCRIPTION OF INVENTION

15 An open reading frame (ORF) encoding the chlamydial lpxB protein has been identified from the *C. pneumoniae* genome. The gene encoding this protein has been inserted into an expression plasmid and shown to confer immune protection against chlamydial infection. Accordingly, this lpxB protein
20 and related polypeptides can be used to prevent and treat *Chlamydia* infection.

 According to a first aspect of the invention, isolated polynucleotides are provided which encode *Chlamydia* polypeptides, whose amino acid sequences are shown in SEQ ID
25 No: 2.

 The term "isolated polynucleotide" is defined as a polynucleotide removed from the environment in which it naturally occurs. For example, a naturally-occurring DNA molecule present in the genome of a living bacteria or as part
30 of a gene bank is not isolated, but the same molecule separated from the remaining part of the bacterial genome, as a result of, e.g., a cloning event (amplification), is isolated. Typically, an isolated DNA molecule is free from DNA regions

(e.g., coding regions) with which it is immediately contiguous at the 5' or 3' end, in the naturally occurring genome. Such isolated polynucleotides may be part of a vector or a composition and still be defined as isolated in that such a vector or composition is not part of the natural environment of such polynucleotide.

The polynucleotide of the invention is either RNA or DNA (cDNA, genomic DNA, or synthetic DNA), or modifications, variants, homologs or fragments thereof. The DNA is either double-stranded or single-stranded, and, if single-stranded, is either the coding strand or the non-coding (anti-sense) strand. Any one of the sequences that encode the polypeptides of the invention as shown in SEQ ID No: 1 is (a) a coding sequence, (b) a ribonucleotide sequence derived from transcription of (a), or (c) a coding sequence which uses the redundancy or degeneracy of the genetic code to encode the same polypeptides. By "polypeptide" or "protein" is meant any chain of amino acids, regardless of length or post-translational modification (e.g., glycosylation or phosphorylation). Both terms are used interchangeably in the present application.

Consistent with the first aspect of the invention, amino acid sequences are provided which are homologous to SEQ ID No: 2. As used herein, "homologous amino acid sequence" is any polypeptide which is encoded, in whole or in part, by a nucleic acid sequence which hybridizes at 25-35°C below critical melting temperature (T_m), to any portion of the nucleic acid sequence of SEQ ID No: 1. A homologous amino acid sequence is one that differs from an amino acid sequence shown in SEQ ID No: 2 by one or more conservative amino acid substitutions. Such a sequence also encompass serotypic variants (defined below) as well as sequences containing deletions or insertions which retain inherent characteristics of the polypeptide such as immunogenicity. Preferably, such a sequence is at least

75%, more preferably 80%, and most preferably 90% identical to SEQ ID No: 2.

Homologous amino acid sequences include sequences that are identical or substantially identical to SEQ ID No: 2.

5 By "amino acid sequence substantially identical" is meant a sequence that is at least 90%, preferably 95%, more preferably 97%, and most preferably 99% identical to an amino acid sequence of reference and that preferably differs from the sequence of reference by a majority of conservative amino acid
10 substitutions.

Conservative amino acid substitutions are substitutions among amino acids of the same class. These classes include, for example, amino acids having uncharged polar side chains, such as asparagine, glutamine, serine,
15 threonine, and tyrosine; amino acids having basic side chains, such as lysine, arginine, and histidine; amino acids having acidic side chains, such as aspartic acid and glutamic acid; and amino acids having nonpolar side chains, such as glycine, alanine, valine, leucine, isoleucine, proline, phenylalanine,
20 methionine, tryptophan, and cysteine.

Homology is measured using sequence analysis software such as Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705. Amino acid
25 sequences are aligned to maximize identity. Gaps may be artificially introduced into the sequence to attain proper alignment. Once the optimal alignment has been set up, the degree of homology is established by recording all of the positions in which the amino acids of both sequences are
30 identical, relative to the total number of positions.

Homologous polynucleotide sequences are defined in a similar way. Preferably, a homologous sequence is one that is at least 45%, more preferably 60%, and most preferably 85% identical to the coding sequence of SEQ ID No: 1.

Consistent with the first aspect of the invention, polypeptides having a sequence homologous to SEQ ID No: 2 include naturally-occurring allelic variants, as well as mutants or any other non-naturally occurring variants that retain the inherent characteristics of the polypeptide of SEQ ID No: 2.

As is known in the art, an allelic variant is an alternate form of a polypeptide that is characterized as having a substitution, deletion, or addition of one or more amino acids that does not alter the biological function of the polypeptide. By "biological function" is meant the function of the polypeptide in the cells in which it naturally occurs, even if the function is not necessary for the growth or survival of the cells. For example, the biological function of a porin is to allow the entry into cells of compounds present in the extracellular medium. Biological function is distinct from antigenic property. A polypeptide can have more than one biological function.

Allelic variants are very common in nature. For example, a bacterial species such as *C. pneumoniae*, is usually represented by a variety of strains that differ from each other by minor allelic variations. Indeed, a polypeptide that fulfills the same biological function in different strains can have an amino acid sequence (and polynucleotide sequence) that is not identical in each of the strains. Despite this variation, an immune response directed generally against many allelic variants has been demonstrated. In studies of the *Chlamydial* MOMP antigen, cross-strain antibody binding plus neutralization of infectivity occurs despite amino acid sequence variation of MOMP from strain to strain, indicating that the MOMP, when used as an immunogen, is tolerant of amino acid variations.

Polynucleotides encoding homologous polypeptides or allelic variants are retrieved by polymerase chain reaction

(PCR) amplification of genomic bacterial DNA extracted by conventional methods. This involves the use of synthetic oligonucleotide primers matching upstream and downstream of the 5' and 3' ends of the encoding domain. Suitable primers are
5 designed according to the nucleotide sequence information provided in SEQ ID No:1. The procedure is as follows: a primer is selected which consists of 10 to 40, preferably 15 to 25 nucleotides. It is advantageous to select primers containing C and G nucleotides in a proportion sufficient to ensure
10 efficient hybridization; *i.e.*, an amount of C and G nucleotides of at least 40%, preferably 50% of the total nucleotide content. A standard PCR reaction contains typically 0.5 to 5 Units of Taq DNA polymerase per 100 μ L, 20 to 200 μ M deoxynucleotide each, preferably at equivalent concentrations,
15 0.5 to 2.5 mM magnesium over the total deoxynucleotide concentration, 10^5 to 10^6 target molecules, and about 20 pmol of each primer. About 25 to 50 PCR cycles are performed, with an annealing temperature 15°C to 5°C below the true T_m of the primers. A more stringent annealing temperature improves
20 discrimination against incorrectly annealed primers and reduces incorporation of incorrect nucleotides at the 3' end of primers. A denaturation temperature of 95°C to 97°C is typical, although higher temperatures may be appropriate for dematuration of G+C-rich targets. The number of cycles performed depends on the
25 starting concentration of target molecules, though typically more than 40 cycles is not recommended as non-specific background products tend to accumulate.

An alternative method for retrieving polynucleotides encoding homologous polypeptides or allelic variants is by
30 hybridization screening of a DNA or RNA library. Hybridization procedures are well-known in the art and are described in Ausubel *et al.*, (Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley & Sons Inc., 1994), Silhavy *et al.* (Silhavy

et al. Experiments with Gene Fusions, Cold Spring Harbor Laboratory Press, 1984), and Davis et al. (Davis et al. A Manual for Genetic Engineering: Advanced Bacterial Genetics, Cold Spring Harbor Laboratory Press, 1980)). Important
5 parameters for optimizing hybridization conditions are reflected in a formula used to obtain the critical melting temperature above which two complementary DNA strands separate from each other (Casey & Davidson, Nucl. Acid Res. (1977) 4:1539). For polynucleotides of about 600 nucleotides or
10 larger, this formula is as follows: $T_m = 81.5 + 0.41 \times (\% \text{ G+C}) + 16.6 \log (\text{cation ion concentration}) - 0.63 \times (\% \text{ formamide}) - 600/\text{base number}$. Under appropriate stringency conditions, hybridization temperature (T_h) is approximately 20 to 40°C, 20 to 25°C, or, preferably 30 to 40°C below the calculated T_m .
15 Those skilled in the art will understand that optimal temperature and salt conditions can be readily determined.

For the polynucleotides of the invention, stringent conditions are achieved for both pre-hybridizing and hybridizing incubations (i) within 4-16 hours at 42°C, in 6 x
20 SSC containing 50% formamide, or (ii) within 4-16 hours at 65°C in an aqueous 6 x SSC solution (1 M NaCl, 0.1 M sodium citrate (pH 7.0)). Typically, hybridization experiments are performed at a temperature from 60 to 68°C, e.g. 65°C. At such a temperature, stringent hybridization conditions can be achieved
25 in 6xSSC, preferably in 2xSSC or 1xSSC, more preferably in 0.5xSSC, 0.3xSSC or 0.1xSSC (in the absence of formamide). 1xSSC contains 0.15 M NaCl and 0.015 M sodium citrate.

Useful homologs and fragments thereof that do not occur naturally are designed using known methods for
30 identifying regions of an antigen that are likely to tolerate amino acid sequence changes and/or deletions. As an example, homologous polypeptides from different species are compared; conserved sequences are identified. The more divergent sequences are the most likely to tolerate sequence changes.

Homology among sequences may be analyzed using, as an example, the BLAST homology searching algorithm of Altschul et al., Nucleic Acids Res.; 25:3389-3402 (1997). Alternatively, sequences are modified such that they become more reactive to
5 T- and/or B-cells, based on computer-assisted analysis of probable T- or B-cell epitopes. Yet another alternative is to mutate a particular amino acid residue or sequence within the polypeptide *in vitro*, then screen the mutant polypeptides for their ability to prevent or treat Chlamydia infection according
10 to the method outlined below.

A person skilled in the art will readily understand that by following the screening process of this invention, it will be determined without undue experimentation whether a particular homolog of SEQ ID No. 2 may be useful in the
15 prevention or treatment of Chlamydia infection. The screening procedure comprises the steps:
immunizing an animal, preferably mouse, with the test homolog or fragment;
inoculating the immunized animal with Chlamydia; and
20 selecting those homologs or fragments which confer protection against Chlamydia.

By "conferring protection" is meant that there is a reduction in severity of any of the effects of Chlamydia infection, in comparison with a control animal which was not
25 immunized with the test homolog or fragment.

Consistent with the first aspect of the invention, polypeptide derivatives are provided that are partial sequences of SEQ ID No. 2, partial sequences of polypeptide sequences homologous to SEQ ID No. 2, polypeptides derived from full-
30 length polypeptides by internal deletion, and fusion proteins.

It is an accepted practice in the field of immunology to use fragments and variants of protein immunogens as vaccines, as all that is required to induce an immune response to a protein is a small (e.g., 8 to 10 amino acid) immunogenic

region of the protein. Various short synthetic peptides corresponding to surface-exposed antigens of pathogens other than *Chlamydia* have been shown to be effective vaccine antigens against their respective pathogens, e.g. an 11 residue peptide
5 of murine mammary tumor virus (Casey & Davidson, Nucl. Acid Res. (1977) 4:1539), a 16-residue peptide of Semliki Forest virus (Snijders et al., 1991. J. Gen. Virol. 72:557-565), and two overlapping peptides of 15 residues each from canine parvovirus (Langeveld et al., Vaccine 12(15):1473-1480, 1994).

10 Accordingly, it will be readily apparent to one skilled in the art, having read the present description, that partial sequences of SEQ ID No: 2 or their homologous amino acid sequences are inherent to the full-length sequences and are taught by the present invention. Such polypeptide
15 fragments preferably are at least 12 amino acids in length. Advantageously, they are at least 20 amino acids, preferably at least 50 amino acids, more preferably at least 75 amino acids, and most preferably at least 100 amino acids in length.

Polynucleotides of 30 to 600 nucleotides encoding
20 partial sequences of sequences homologous to SEQ ID No: 2 are retrieved by PCR amplification using the parameters outlined above and using primers matching the sequences upstream and downstream of the 5' and 3' ends of the fragment to be amplified. The template polynucleotide for such amplification
25 is either the full length polynucleotide homologous to SEQ ID No: 1, or a polynucleotide contained in a mixture of polynucleotides such as a DNA or RNA library. As an alternative method for retrieving the partial sequences, screening hybridization is carried out under conditions
30 described above and using the formula for calculating T_m . Where fragments of 30 to 600 nucleotides are to be retrieved, the calculated T_m is corrected by subtracting $(600/\text{polynucleotide size in base pairs})$ and the stringency conditions are defined by a hybridization temperature that is

5 to 10°C below T_m. Where oligonucleotides shorter than 20-30 bases are to be obtained, the formula for calculating the T_m is as follows: $T_m = 4 \times (G+C) + 2 (A+T)$. For example, an 18 nucleotide fragment of 50% G+C would have an approximate T_m of 54°C. Short peptides that are fragments of SEQ ID No: 2 or its homologous sequences, are obtained directly by chemical synthesis (E. Gross and H. J. Meinhofer, 4 The Peptides: Analysis, Synthesis, Biology; Modern Techniques of Peptide Synthesis, John Wiley & Sons (1981), and M. Bodanzki, Principles of Peptide Synthesis, Springer -Verlag (1984)).

Useful polypeptide derivatives, e.g., polypeptide fragments, are designed using computer-assisted analysis of amino acid sequences. This would identify probable surface-exposed, antigenic regions (Hughes et al., 1992. Infect. Immun. 60(9):3497). Analysis of 6 amino acid sequences contained in SEQ ID No: 2, based on the product of flexibility and hydrophobicity propensities using the program SEQSEE (Wishart DS, et al. "SEQSEE: a comprehensive program suite for protein sequence analysis." *Comput Appl Biosci*. 1994 Apr;10(2):121-32), can reveal potential B- and T-cell epitopes which may be used as a basis for selecting useful immunogenic fragments and variants. This analysis uses a reasonable combination of external surface features that is likely to be recognized by antibodies. Probable T-cell epitopes for HLA-A0201 MHC subclass may be revealed by an algorithms that emulate an approach developed at the NIH (Parker KC, et al. "Peptide binding to MHC class I molecules: implications for antigenic peptide prediction." *Immunol Res* 1995;14(1):34-57).

Epitopes which induce a protective T cell-dependent immune response are present throughout the length of the polypeptide. However, some epitopes may be masked by secondary and tertiary structures of the polypeptide. To reveal such masked epitopes large internal deletions are created which remove much of the original protein structure and exposes the

masked epitopes. Such internal deletions sometimes effect the additional advantage of removing immunodominant regions of high variability among strains.

Polynucleotides encoding polypeptide fragments and
5 polypeptides having large internal deletions are constructed using standard methods (Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley & Sons Inc., 1994). Such methods include standard PCR, inverse PCR, restriction enzyme treatment of cloned DNA molecules, or the method of Kunkel *et al.*
10 (Kunkel *et al.* Proc. Natl. Acad. Sci. USA (1985) 82:448). Components for these methods and instructions for their use are readily available from various commercial sources such as Stratagene. Once the deletion mutants have been constructed, they are tested for their ability to prevent or treat Chlamydia
15 infection as described above.

As used herein, a fusion polypeptide is one that contains a polypeptide or a polypeptide derivative of the invention fused at the N- or C-terminal end to any other polypeptide (hereinafter referred to as a peptide tail). A
20 simple way to obtain such a fusion polypeptide is by translation of an in-frame fusion of the polynucleotide sequences, *i.e.*, a hybrid gene. The hybrid gene encoding the fusion polypeptide is inserted into an expression vector which is used to transform or transfect a host cell. Alternatively,
25 the polynucleotide sequence encoding the polypeptide or polypeptide derivative is inserted into an expression vector in which the polynucleotide encoding the peptide tail is already present. Such vectors and instructions for their use are commercially available, *e.g.* the pMal-c2 or pMal-p2 system from
30 New England Biolabs, in which the peptide tail is a maltose binding protein, the glutathione-S-transferase system of Pharmacia, or the His-Tag system available from Novagen. These and other expression systems provide convenient means for

further purification of polypeptides and derivatives of the invention.

An advantageous example of a fusion polypeptide is one where the polypeptide or homolog or fragment of the invention is fused to a polypeptide having adjuvant activity, such as subunit B of either cholera toxin or *E. coli* heat-labile toxin. Another advantageous fusion is one where the polypeptide, homolog or fragment is fused to a strong T-cell epitope or B-cell epitope. Such an epitope may be one known in the art (e.g. the Hepatitis B virus core antigen, D.R. Millich et al., "Antibody production to the nucleocapsid and envelope of the Hepatitis B virus primed by a single synthetic T cell site", *Nature*. 1987. 329:547-549), or one which has been identified in another polypeptide of the invention based on computer-assisted analysis of probable T- or B-cell epitopes. Consistent with this aspect of the invention is a fusion polypeptide comprising T- or B-cell epitopes from SEQ ID No: 2 or its homolog or fragment, wherein the epitopes are derived from multiple variants of said polypeptide or homolog or fragment, each variant differing from another in the location and sequence of its epitope within the polypeptide. Such a fusion is effective in the prevention and treatment of Chlamydia infection since it optimizes the T- and B-cell response to the overall polypeptide, homolog or fragment.

To effect fusion, the polypeptide of the invention is fused to the N-, or preferably, to the C-terminal end of the polypeptide having adjuvant activity or T- or B-cell epitope. Alternatively, a polypeptide fragment of the invention is inserted internally within the amino acid sequence of the polypeptide having adjuvant activity. The T- or B-cell epitope may also be inserted internally within the amino acid sequence of the polypeptide of the invention.

Consistent with the first aspect, the polynucleotides of the invention also encode hybrid precursor polypeptides

containing heterologous signal peptides, which mature into polypeptides of the invention. By "heterologous signal peptide" is meant a signal peptide that is not found in naturally-occurring precursors of polypeptides of the invention.

Polynucleotide molecules according to the invention, including RNA, DNA, or modifications or combinations thereof, have various applications. A DNA molecule is used, for example, (i) in a process for producing the encoded polypeptide in a recombinant host system, (ii) in the construction of vaccine vectors such as poxviruses, which are further used in methods and compositions for preventing and/or treating *Chlamydia* infection, (iii) as a vaccine agent (as well as an RNA molecule), in a naked form or formulated with a delivery vehicle and, (iv) in the construction of attenuated *Chlamydia* strains that can over-express a polynucleotide of the invention or express it in a non-toxic, mutated form.

Accordingly, a second aspect of the invention encompasses (i) an expression cassette containing a DNA molecule of the invention placed under the control of the elements required for expression, in particular under the control of an appropriate promoter; (ii) an expression vector containing an expression cassette of the invention; (iii) a procaryotic or eucaryotic cell transformed or transfected with an expression cassette and/or vector of the invention, as well as (iv) a process for producing a polypeptide or polypeptide derivative encoded by a polynucleotide of the invention, which involves culturing a procaryotic or eucaryotic cell transformed or transfected with an expression cassette and/or vector of the invention, under conditions that allow expression of the DNA molecule of the invention and, recovering the encoded polypeptide or polypeptide derivative from the cell culture.

A recombinant expression system is selected from procaryotic and eucaryotic hosts. Eucaryotic hosts include

yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris*), mammalian cells (e.g., COS1, NIH3T3, or JEG3 cells), arthropods cells (e.g., *Spodoptera frugiperda* (SF9) cells), and plant cells. A preferred expression system is a procaryotic
5 host such as *E. coli*. Bacterial and eucaryotic cells are available from a number of different sources including commercial sources to those skilled in the art, e.g., the American Type Culture Collection (ATCC; Rockville, Maryland). Commercial sources of cells used for recombinant protein
10 expression also provide instructions for usage of the cells.

The choice of the expression system depends on the features desired for the expressed polypeptide. For example, it may be useful to produce a polypeptide of the invention in a particular lipidated form or any other form.

15 One skilled in the art would readily understand that not all vectors and expression control sequences and hosts would be expected to express equally well the polynucleotides of this invention. With the guidelines described below, however, a selection of vectors, expression control sequences
20 and hosts may be made without undue experimentation and without departing from the scope of this invention.

In selecting a vector, the host must be chosen that is compatible with the vector which is to exist and possibly replicate in it. Considerations are made with respect to the
25 vector copy number, the ability to control the copy number, expression of other proteins such as antibiotic resistance. In selecting an expression control sequence, a number of variables are considered. Among the important variable are the relative strength of the sequence (e.g. the ability to drive expression
30 under various conditions), the ability to control the sequence's function, compatibility between the polynucleotide to be expressed and the control sequence (e.g. secondary structures are considered to avoid hairpin structures which prevent efficient transcription). In selecting the host,

unicellular hosts are selected which are compatible with the selected vector, tolerant of any possible toxic effects of the expressed product, able to secrete the expressed product efficiently if such is desired, to be able to express the product in the desired conformation, to be easily scaled up, and to which ease of purification of the final product.

The choice of the expression cassette depends on the host system selected as well as the features desired for the expressed polypeptide. Typically, an expression cassette includes a promoter that is functional in the selected host system and can be constitutive or inducible; a ribosome binding site; a start codon (ATG) if necessary; a region encoding a signal peptide, e.g., a lipidation signal peptide; a DNA molecule of the invention; a stop codon; and optionally a 3' terminal region (translation and/or transcription terminator). The signal peptide encoding region is adjacent to the polynucleotide of the invention and placed in proper reading frame. The signal peptide-encoding region is homologous or heterologous to the DNA molecule encoding the mature polypeptide and is compatible with the secretion apparatus of the host used for expression. The open reading frame constituted by the DNA molecule of the invention, solely or together with the signal peptide, is placed under the control of the promoter so that transcription and translation occur in the host system. Promoters and signal peptide encoding regions are widely known and available to those skilled in the art and include, for example, the promoter of *Salmonella typhimurium* (and derivatives) that is inducible by arabinose (promoter araB) and is functional in Gram-negative bacteria such as *E. coli* (as described in U.S. Patent No. 5,028,530 and in Cagnon et al., (Cagnon et al., Protein Engineering (1991) 4(7):843)); the promoter of the gene of bacteriophage T7 encoding RNA polymerase, that is functional in a number of *E. coli* strains expressing T7 polymerase (described in U.S. Patent

No. 4,952,496); OspA lipidation signal peptide ; and RlpB lipidation signal peptide (Takase et al., J. Bact. (1987) 169:5692).

The expression cassette is typically part of an
5 expression vector, which is selected for its ability to replicate in the chosen expression system. Expression vectors (e.g., plasmids or viral vectors) can be chosen, for example, from those described in Pouwels et al. (Cloning Vectors: A Laboratory Manual 1985, Supp. 1987). Suitable expression
10 vectors can be purchased from various commercial sources.

Methods for transforming/transfecting host cells with expression vectors are well-known in the art and depend on the host system selected as described in Ausubel et al., (Ausubel et al., Current Protocols in Molecular Biology, John Wiley &
15 Sons Inc., 1994).

Upon expression, a recombinant polypeptide of the invention (or a polypeptide derivative) is produced and remains in the intracellular compartment, is secreted/excreted in the extracellular medium or in the periplasmic space, or is
20 embedded in the cellular membrane. The polypeptide is recovered in a substantially purified form from the cell extract or from the supernatant after centrifugation of the recombinant cell culture. Typically, the recombinant polypeptide is purified by antibody-based affinity purification
25 or by other well-known methods that can be readily adapted by a person skilled in the art, such as fusion of the polynucleotide encoding the polypeptide or its derivative to a small affinity binding domain. Antibodies useful for purifying by immunoaffinity the polypeptides of the invention are obtained
30 as described below.

A polynucleotide of the invention can also be useful as a vaccine. There are two major routes, either using a viral or bacterial host as gene delivery vehicle (live vaccine vector) or administering the gene in a free form, e.g.,

inserted into a plasmid. Therapeutic or prophylactic efficacy of a polynucleotide of the invention is evaluated as described below.

Accordingly, a third aspect of the invention provides

5 (i) a vaccine vector such as a poxvirus, containing a DNA molecule of the invention, placed under the control of elements required for expression; (ii) a composition of matter comprising a vaccine vector of the invention, together with a diluent or carrier; specifically (iii) a pharmaceutical

10 composition containing a therapeutically or prophylactically effective amount of a vaccine vector of the invention; (iv) a method for inducing an immune response against *Chlamydia* in a mammal (e.g., a human; alternatively, the method can be used in veterinary applications for treating or preventing *Chlamydia*

15 infection of animals, e.g., cats or birds), which involves administering to the mammal an immunogenically effective amount of a vaccine vector of the invention to elicit a protective or therapeutic immune response to *Chlamydia* ; and particularly, (v) a method for preventing and/or treating a *Chlamydia* (e.g.,

20 *C. trachomatis*, *C. psittaci*, *C. pneumonia*, *C. pecorum*) infection, which involves administering a prophylactic or therapeutic amount of a vaccine vector of the invention to an infected individual. Additionally, the third aspect of the invention encompasses the use of a vaccine vector of the

25 invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection.

As used herein, a vaccine vector expresses one or several polypeptides or derivatives of the invention. The vaccine vector may express additionally a cytokine, such as

30 interleukin-2 (IL-2) or interleukin-12 (IL-12), that enhances the immune response (adjuvant effect). It is understood that each of the components to be expressed is placed under the control of elements required for expression in a mammalian cell.

Consistent with the third aspect of the invention is a composition comprising several vaccine vectors, each of them capable of expressing a polypeptide or derivative of the invention. A composition may also comprise a vaccine vector
5 capable of expressing an additional *Chlamydia* antigen, or a subunit, fragment, homolog, mutant, or derivative thereof; optionally together with or a cytokine such as IL-2 or IL-12.

Vaccination methods for treating or preventing infection in a mammal comprises use of a vaccine vector of the
10 invention to be administered by any conventional route, particularly to a mucosal (e.g., ocular, intranasal, oral, gastric, pulmonary, intestinal, rectal, vaginal, or urinary tract) surface or via the parenteral (e.g., subcutaneous, intradermal, intramuscular, intravenous, or intraperitoneal)
15 route. Preferred routes depend upon the choice of the vaccine vector. Treatment may be effected in a single dose or repeated at intervals. The appropriate dosage depends on various parameters understood by skilled artisans such as the vaccine vector itself, the route of administration or the condition of
20 the mammal to be vaccinated (weight, age and the like).

Live vaccine vectors available in the art include viral vectors such as adenoviruses and poxviruses as well as bacterial vectors, e.g., *Shigella*, *Salmonella*, *Vibrio cholerae*, *Lactobacillus*, Bacille bilié de Calmette-Guérin (BCG), and
25 *Streptococcus*.

An example of an adenovirus vector, as well as a method for constructing an adenovirus vector capable of expressing a DNA molecule of the invention, are described in U.S. Patent No. 4,920,209. Poxvirus vectors include vaccinia
30 and canary pox virus, described in U.S. Patent No. 4,722,848 and U.S. Patent No. 5,364,773, respectively. (Also see, e.g., Tartaglia et al., Virology (1992) 188:217) for a description of a vaccinia virus vector and Taylor et al, Vaccine (1995) 13:539 for a reference of a canary pox.) Poxvirus vectors capable of

expressing a polynucleotide of the invention are obtained by homologous recombination as described in Kieny et al., Nature (1984) 312:163 so that the polynucleotide of the invention is inserted in the viral genome under appropriate conditions for expression in mammalian cells. Generally, the dose of vaccine viral vector, for therapeutic or prophylactic use, can be of from about 1×10^4 to about 1×10^{11} , advantageously from about 1×10^7 to about 1×10^{10} , preferably of from about 1×10^7 to about 1×10^9 plaque-forming units per kilogram. Preferably, viral vectors are administered parenterally; for example, in 3 doses, 4 weeks apart. It is preferable to avoid adding a chemical adjuvant to a composition containing a viral vector of the invention and thereby minimizing the immune response to the viral vector itself.

Non-toxicogenic *Vibrio cholerae* mutant strains that are useful as a live oral vaccine are known. Mekalanos et al., Nature (1983) 306:551 and U.S. Patent No. 4,882,278 describe strains which have a substantial amount of the coding sequence of each of the two *ctxA* alleles deleted so that no functional *cholerae* toxin is produced. WO 92/11354 describes a strain in which the *irgA* locus is inactivated by mutation; this mutation can be combined in a single strain with *ctxA* mutations. WO 94/01533 describes a deletion mutant lacking functional *ctxA* and *attRS1* DNA sequences. These mutant strains are genetically engineered to express heterologous antigens, as described in WO 94/19482. An effective vaccine dose of a *Vibrio cholerae* strain capable of expressing a polypeptide or polypeptide derivative encoded by a DNA molecule of the invention contains about 1×10^5 to about 1×10^9 , preferably about 1×10^6 to about 1×10^8 , viable bacteria in a volume appropriate for the selected route of administration. Preferred routes of administration include all mucosal routes; most preferably, these vectors are administered intranasally or orally.

Attenuated *Salmonella typhimurium* strains, genetically engineered for recombinant expression of heterologous antigens or not, and their use as oral vaccines are described in Nakayama et al. (Bio/Technology (1988) 6:693) and WO 92/11361. Preferred routes of administration include all mucosal routes; most preferably, these vectors are administered intranasally or orally.

Other bacterial strains used as vaccine vectors in the context of the present invention are described for *Shigella flexneri* in High et al., EMBO (1992) 11:1991 and Sizemore et al., Science (1995) 270:299; for *Streptococcus gordonii* in Medaglini et al., Proc. Natl. Acad. Sci. USA (1995) 92:6868; and for Bacille Calmette Guerin in Flynn J.L., Cell. Mol. Biol. (1994) 40 (suppl. I):31, WO 88/06626, WO 90/00594, WO 91/13157, WO 92/01796, and WO 92/21376.

In bacterial vectors, the polynucleotide of the invention is inserted into the bacterial genome or remains in a free state as part of a plasmid.

The composition comprising a vaccine bacterial vector of the present invention may further contain an adjuvant. A number of adjuvants are known to those skilled in the art. Preferred adjuvants are selected as provided below.

Accordingly, a fourth aspect of the invention provides (i) a composition of matter comprising a polynucleotide of the invention, together with a diluent or carrier; (ii) a pharmaceutical composition comprising a therapeutically or prophylactically effective amount of a polynucleotide of the invention; (iii) a method for inducing an immune response against *Chlamydia* in a mammal by administration of an immunogenically effective amount of a polynucleotide of the invention to elicit a protective immune response to *Chlamydia*; and particularly, (iv) a method for preventing and/or treating a *Chlamydia* (e.g., *C. trachomatis*, *C. psittaci*, *C. pneumoniae*, or *C. pecorum*) infection, by administering a

prophylactic or therapeutic amount of a polynucleotide of the invention to an infected individual. Additionally, the fourth aspect of the invention encompasses the use of a polynucleotide of the invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection. A preferred use includes the use of a DNA molecule placed under conditions for expression in a mammalian cell, especially in a plasmid that is unable to replicate in mammalian cells and to substantially integrate in a mammalian genome.

Use of the polynucleotides of the invention include their administration to a mammal as a vaccine, for therapeutic or prophylactic purposes. Such polynucleotides are used in the form of DNA as part of a plasmid that is unable to replicate in a mammalian cell and unable to integrate into the mammalian genome. Typically, such a DNA molecule is placed under the control of a promoter suitable for expression in a mammalian cell. The promoter functions either ubiquitously or tissue-specifically. Examples of non-tissue specific promoters include the early Cytomegalovirus (CMV) promoter (described in U.S. Patent No. 4,168,062) and the Rous Sarcoma Virus promoter (described in Norton & Coffin, *Molec. Cell Biol.* (1985) 5:281). An example of a tissue-specific promoter is the desmin promoter which drives expression in muscle cells (Li *et al.*, *Gene* (1989) 78:243, Li & Paulin, *J. Biol. Chem.* (1991) 266:6562 and Li & Paulin, *J. Biol. Chem.* (1993) 268:10403). Use of promoters is well-known to those skilled in the art. Useful vectors are described in numerous publications, specifically WO 94/21797 and Hartikka *et al.*, *Human Gene Therapy* (1996) 7:1205.

Polynucleotides of the invention which are used as vaccines encode either a precursor or a mature form of the corresponding polypeptide. In the precursor form, the signal peptide is either homologous or heterologous. In the latter case, a eucaryotic leader sequence such as the leader sequence of the tissue-type plasminogen factor (tPA) is preferred.

As used herein, a composition of the invention contains one or several polynucleotides with optionally at least one additional polynucleotide encoding another *Chlamydia* antigen such as urease subunit A, B, or both, or a fragment, derivative, mutant, or analog thereof. The composition may also contain an additional polynucleotide encoding a cytokine, such as interleukin-2 (IL-2) or interleukin-12 (IL-12) so that the immune response is enhanced. These additional polynucleotides are placed under appropriate control for expression. Advantageously, DNA molecules of the invention and/or additional DNA molecules to be included in the same composition, are present in the same plasmid.

Standard techniques of molecular biology for preparing and purifying polynucleotides are used in the preparation of polynucleotide therapeutics of the invention. For use as a vaccine, a polynucleotide of the invention is formulated according to various methods outlined below.

One method utilizes the polynucleotide in a naked form, free of any delivery vehicles. Such a polynucleotide is simply diluted in a physiologically acceptable solution such as sterile saline or sterile buffered saline, with or without a carrier. When present, the carrier preferably is isotonic, hypotonic, or weakly hypertonic, and has a relatively low ionic strength, such as provided by a sucrose solution, e.g., a solution containing 20% sucrose.

An alternative method utilizes the polynucleotide in association with agents that assist in cellular uptake. Examples of such agents are (i) chemicals that modify cellular permeability, such as bupivacaine (see, e.g., WO 94/16737), (ii) liposomes for encapsulation of the polynucleotide, or (iii) cationic lipids or silica, gold, or tungsten microparticles which associate themselves with the polynucleotides.

Anionic and neutral liposomes are well-known in the art (see, e.g., Liposomes: A Practical Approach, RPC New Ed, IRL press (1990), for a detailed description of methods for making liposomes) and are useful for delivering a large range
5 of products, including polynucleotides.

Cationic lipids are also known in the art and are commonly used for gene delivery. Such lipids include LipofectinTM also known as DOTMA (N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride), DOTAP (1,2-bis(oleyloxy)-3-
10 (trimethylammonio)propane), DDAB (dimethyldioctadecylammonium bromide), DOGS (dioctadecylamidologlycyl spermine) and cholesterol derivatives such as DC-Chol (3 beta-(N-(N',N'-dimethyl aminomethane)-carbamoyl) cholesterol). A description of these cationic lipids can be found in EP 187,702,
15 WO 90/11092, U.S. Patent No. 5,283,185, WO 91/15501, WO 95/26356, and U.S. Patent No. 5,527,928. Cationic lipids for gene delivery are preferably used in association with a neutral lipid such as DOPE (dioleoyl phosphatidylethanolamine), as described in WO 90/11092 as an example.

20 Formulation containing cationic liposomes may optionally contain other transfection-facilitating compounds. A number of them are described in WO 93/18759, WO 93/19768, WO 94/25608, and WO 95/02397. They include spermine derivatives useful for facilitating the transport of DNA through the
25 nuclear membrane (see, for example, WO 93/18759) and membrane-permeabilizing compounds such as GALA, Gramicidine S, and cationic bile salts (see, for example, WO 93/19768).

Gold or tungsten microparticles are used for gene delivery, as described in WO 91/00359, WO 93/17706, and Tang et
30 al. Nature (1992) 356:152. The microparticle-coated polynucleotide is injected via intradermal or intraepidermal routes using a needleless injection device ("gene gun"), such as those described in U.S. Patent No. 4,945,050, U.S. Patent No. 5,015,580, and WO 94/24263.

The amount of DNA to be used in a vaccine recipient depends, e.g., on the strength of the promoter used in the DNA construct, the immunogenicity of the expressed gene product, the condition of the mammal intended for administration (e.g.,
5 the weight, age, and general health of the mammal), the mode of administration, and the type of formulation. In general, a therapeutically or prophylactically effective dose from about 1 μ g to about 1 mg, preferably, from about 10 μ g to about 800 μ g and, more preferably, from about 25 μ g to about 250 μ g, can
10 be administered to human adults. The administration can be achieved in a single dose or repeated at intervals.

The route of administration is any conventional route used in the vaccine field. As general guidance, a polynucleotide of the invention is administered *via* a mucosal
15 surface, e.g., an ocular, intranasal, pulmonary, oral, intestinal, rectal, vaginal, and urinary tract surface; or *via* a parenteral route, e.g., by an intravenous, subcutaneous, intraperitoneal, intradermal, intraepidermal, or intramuscular route. The choice of administration route depends on the
20 formulation that is selected. A polynucleotide formulated in association with bupivacaine is advantageously administered into muscles. When a neutral or anionic liposome or a cationic lipid, such as DOTMA or DC-Chol, is used, the formulation can be advantageously injected *via* intravenous, intranasal
25 (aerosolization), intramuscular, intradermal, and subcutaneous routes. A polynucleotide in a naked form can advantageously be administered *via* the intramuscular, intradermal, or subcutaneous routes.

Although not absolutely required, such a composition
30 can also contain an adjuvant. If so, a systemic adjuvant that does not require concomitant administration in order to exhibit an adjuvant effect is preferable such as, e.g., QS21, which is described in U.S. Patent No. 5,057,546.

The sequence information provided in the present application enables the design of specific nucleotide probes and primers that are used for diagnostic purposes.

Accordingly, a fifth aspect of the invention provides a
5 nucleotide probe or primer having a sequence found in or derived by degeneracy of the genetic code from a sequence shown in SEQ ID No:1.

The term "probe" as used in the present application refers to DNA (preferably single stranded) or RNA molecules (or
10 modifications or combinations thereof) that hybridize under the stringent conditions, as defined above, to nucleic acid molecules having SEQ ID No:1 or to sequences homologous to SEQ ID No:1, or to its complementary or anti-sense sequence. Generally, probes are significantly shorter than full-length
15 sequences. Such probes contain from about 5 to about 100, preferably from about 10 to about 80, nucleotides. In particular, probes have sequences that are at least 75%, preferably at least 85%, more preferably 95% homologous to a portion of SEQ ID No:1 or that are complementary to such
20 sequences. Probes may contain modified bases such as inosine, methyl-5-deoxycytidine, deoxyuridine, dimethylamino-5-deoxyuridine, or diamino-2, 6-purine. Sugar or phosphate residues may also be modified or substituted. For example, a deoxyribose residue may be replaced by a polyamide (Nielsen
25 et al., Science (1991) 254:1497) and phosphate residues may be replaced by ester groups such as diphosphate, alkyl, arylphosphonate and phosphorothioate esters. In addition, the 2'-hydroxyl group on ribonucleotides may be modified by including such groups as alkyl groups.

30 Probes of the invention are used in diagnostic tests, as capture or detection probes. Such capture probes are conventionally immobilized on a solid support, directly or indirectly, by covalent means or by passive adsorption. A detection probe is labelled by a detection marker selected

from: radioactive isotopes, enzymes such as peroxidase, alkaline phosphatase, and enzymes able to hydrolyze a chromogenic, fluorogenic, or luminescent substrate, compounds that are chromogenic, fluorogenic, or luminescent, nucleotide
5 base analogs, and biotin.

Probes of the invention are used in any conventional hybridization technique, such as dot blot (Maniatis *et al.*, Molecular Cloning: A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York),
10 Southern blot (Southern, J. Mol. Biol. (1975) 98:503), northern blot (identical to Southern blot with the exception that RNA is used as a target), or the sandwich technique (Dunn *et al.*, Cell (1977) 12:23). The latter technique involves the use of a specific capture probe and/or a specific detection probe with
15 nucleotide sequences that at least partially differ from each other.

A primer is a probe of usually about 10 to about 40 nucleotides that is used to initiate enzymatic polymerization of DNA in an amplification process (e.g., PCR),
20 in an elongation process, or in a reverse transcription method. Primers used in diagnostic methods involving PCR are labeled by methods known in the art.

As described herein, the invention also encompasses (i) a reagent comprising a probe of the invention for detecting
25 and/or identifying the presence of *Chlamydia* in a biological material; (ii) a method for detecting and/or identifying the presence of *Chlamydia* in a biological material, in which (a) a sample is recovered or derived from the biological material, (b) DNA or RNA is extracted from the material and denatured,
30 and (c) exposed to a probe of the invention, for example, a capture, detection probe or both, under stringent hybridization conditions, such that hybridization is detected; and (iii) a method for detecting and/or identifying the presence of *Chlamydia* in a biological material, in which (a) a sample is

recovered or derived from the biological material, (b) DNA is extracted therefrom, (c) the extracted DNA is primed with at least one, and preferably two, primers of the invention and amplified by polymerase chain reaction, and (d) the amplified
5 DNA fragment is produced.

It is apparent that disclosure of polynucleotide sequences of SEQ ID No:1, its homologs and partial sequences enable their corresponding amino acid sequences. Accordingly, a sixth aspect of the invention features a substantially
10 purified polypeptide or polypeptide derivative having an amino acid sequence encoded by a polynucleotide of the invention.

A "substantially purified polypeptide" as used herein is defined as a polypeptide that is separated from the environment in which it naturally occurs and/or that is free of
15 the majority of the polypeptides that are present in the environment in which it was synthesized. For example, a substantially purified polypeptide is free from cytoplasmic polypeptides. Those skilled in the art would readily understand that the polypeptides of the invention may be
20 purified from a natural source, *i.e.*, a *Chlamydia* strain, or produced by recombinant means.

Consistent with the sixth aspect of the invention are polypeptides, homologs or fragments which are modified or treated to enhance their immunogenicity in the target animal,
25 in whom the polypeptide, homolog or fragments are intended to confer protection against *Chlamydia*. Such modifications or treatments include: amino acid substitutions with an amino acid derivative such as 3-methylhistidine, 4-hydroxyproline, 5-hydroxylysine etc., modifications or deletions which are
30 carried out after preparation of the polypeptide, homolog or fragment, such as the modification of free amino, carboxyl or hydroxyl side groups of the amino acids.

Identification of homologous polypeptides or polypeptide derivatives encoded by polynucleotides of the

invention which have specific antigenicity is achieved by screening for cross-reactivity with an antiserum raised against the polypeptide of reference having an amino acid sequence of SEQ ID No:1. The procedure is as follows: a monospecific
5 hyperimmune antiserum is raised against a purified reference polypeptide, a fusion polypeptide (for example, an expression product of MBP, GST, or His-tag systems, the description and instructions for use of which are contained in Invitrogen product manuals for pcDNA3.1/Myc-His(+) A, B, and C and for the
10 Xpress[™] System Protein Purification), or a synthetic peptide predicted to be antigenic. Where an antiserum is raised against a fusion polypeptide, two different fusion systems are employed. Specific antigenicity can be determined according to a number of methods, including Western blot (Towbin et al.,
15 Proc. Natl. Acad. Sci. USA (1979) 76:4350), dot blot, and ELISA, as described below.

In a Western blot assay, the product to be screened, either as a purified preparation or a total *E. coli* extract, is submitted to SDS-Page electrophoresis as described by Laemmli
20 (Nature (1970) 227:680). After transfer to a nitrocellulose membrane, the material is further incubated with the monospecific hyperimmune antiserum diluted in the range of dilutions from about 1:5 to about 1:5000, preferably from about 1:100 to about 1:500. Specific antigenicity is shown once a
25 band corresponding to the product exhibits reactivity at any of the dilutions in the above range.

In an ELISA assay, the product to be screened is preferably used as the coating antigen. A purified preparation is preferred, although a whole cell extract can also be used.
30 Briefly, about 100 μ l of a preparation at about 10 μ g protein/ml are distributed into wells of a 96-well polycarbonate ELISA plate. The plate is incubated for 2 hours at 37°C then overnight at 4°C. The plate is washed with phosphate buffer saline (PBS) containing 0.05% Tween 20

(PBS/Tween buffer). The wells are saturated with 250 μ l PBS containing 1% bovine serum albumin (BSA) to prevent non-specific antibody binding. After 1 hour incubation at 37°C, the plate is washed with PBS/Tween buffer. The antiserum is
5 serially diluted in PBS/Tween buffer containing 0.5% BSA. 100 μ l of dilutions are added per well. The plate is incubated for 90 minutes at 37°C, washed and evaluated according to standard procedures. For example, a goat anti-rabbit peroxidase conjugate is added to the wells when specific antibodies were
10 raised in rabbits. Incubation is carried out for 90 minutes at 37°C and the plate is washed. The reaction is developed with the appropriate substrate and the reaction is measured by colorimetry (absorbance measured spectrophotometrically). Under the above experimental conditions, a positive reaction is
15 shown by O.D. values greater than a non immune control serum.

In a dot blot assay, a purified product is preferred, although a whole cell extract can also be used. Briefly, a solution of the product at about 100 μ g/ml is serially two-fold diluted in 50 mM Tris-HCl (pH 7.5). 100 μ l of each dilution
20 are applied to a nitrocellulose membrane 0.45 μ m set in a 96-well dot blot apparatus (Biorad). The buffer is removed by applying vacuum to the system. Wells are washed by addition of 50 mM Tris-HCl (pH 7.5) and the membrane is air-dried. The membrane is saturated in blocking buffer (50 mM Tris-HCl (pH
25 7.5) 0.15 M NaCl, 10 g/L skim milk) and incubated with an antiserum dilution from about 1:50 to about 1:5000, preferably about 1:500. The reaction is revealed according to standard procedures. For example, a goat anti-rabbit peroxidase conjugate is added to the wells when rabbit antibodies are
30 used. Incubation is carried out 90 minutes at 37°C and the blot is washed. The reaction is developed with the appropriate substrate and stopped. The reaction is measured visually by the appearance of a colored spot, e.g., by colorimetry. Under the above experimental conditions, a positive reaction is shown

once a colored spot is associated with a dilution of at least about 1:5, preferably of at least about 1:500.

Therapeutic or prophylactic efficacy of a polypeptide or derivative of the invention can be evaluated as described below. A seventh aspect of the invention provides (i) a composition of matter comprising a polypeptide of the invention together with a diluent or carrier; specifically (ii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of a polypeptide of the invention; (iii) a method for inducing an immune response against *Chlamydia* in a mammal, by administering to the mammal an immunogenically effective amount of a polypeptide of the invention to elicit a protective immune response to *Chlamydia*; and particularly, (iv) a method for preventing and/or treating a *Chlamydia* (e.g., *C. trachomatis*, *C. psittaci*, *C. pneumoniae*, or *C. pecorum*) infection, by administering a prophylactic or therapeutic amount of a polypeptide of the invention to an infected individual. Additionally, the seventh aspect of the invention encompasses the use of a polypeptide of the invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection.

As used herein, the immunogenic compositions of the invention are administered by conventional routes known in the vaccine field, in particular to a mucosal (e.g., ocular, intranasal, pulmonary, oral, gastric, intestinal, rectal, vaginal, or urinary tract) surface or via the parenteral (e.g., subcutaneous, intradermal, intramuscular, intravenous, or intraperitoneal) route. The choice of administration route depends upon a number of parameters, such as the adjuvant associated with the polypeptide. If a mucosal adjuvant is used, the intranasal or oral route is preferred. If a lipid formulation or an aluminum compound is used, the parenteral route is preferred with the sub-cutaneous or intramuscular route being most preferred. The choice also depends upon the

nature of the vaccine agent. For example, a polypeptide of the invention fused to CTB or LTB is best administered to a mucosal surface.

As used herein, the composition of the invention
5 contains one or several polypeptides or derivatives of the invention. The composition optionally contains at least one additional *Chlamydia* antigen, or a subunit, fragment, homolog, mutant, or derivative thereof.

For use in a composition of the invention, a
10 polypeptide or derivative thereof is formulated into or with liposomes, preferably neutral or anionic liposomes, microspheres, ISCOMS, or virus-like-particles (VLPs) to facilitate delivery and/or enhance the immune response. These compounds are readily available to one skilled in the art; for
15 example, see Liposomes: A Practical Approach, RCP New Ed, IRL press (1990).

Adjuvants other than liposomes and the like are also used and are known in the art. Adjuvants may protect the antigen from rapid dispersal by sequestering it in a local
20 deposit, or they may contain substances that stimulate the host to secrete factors that are chemotactic for macrophages and other components of the immune system. An appropriate selection can conventionally be made by those skilled in the art, for example, from those described below (under the
25 eleventh aspect of the invention).

Treatment is achieved in a single dose or repeated as necessary at intervals, as can be determined readily by one skilled in the art. For example, a priming dose is followed by three booster doses at weekly or monthly intervals. An
30 appropriate dose depends on various parameters including the recipient (e.g., adult or infant), the particular vaccine antigen, the route and frequency of administration, the presence/absence or type of adjuvant, and the desired effect (e.g., protection and/or treatment), as can be determined by

one skilled in the art. In general, a vaccine antigen of the invention is administered by a mucosal route in an amount from about 10 μ g to about 500 mg, preferably from about 1 mg to about 200 mg. For the parenteral route of administration, the
5 dose usually does not exceed about 1 mg, preferably about 100 μ g.

When used as vaccine agents, polynucleotides and polypeptides of the invention may be used sequentially as part of a multistep immunization process. For example, a mammal is
10 initially primed with a vaccine vector of the invention such as a pox virus, e.g., via the parenteral route, and then boosted twice with the polypeptide encoded by the vaccine vector, e.g., via the mucosal route. In another example, liposomes associated with a polypeptide or derivative of the invention is
15 also used for priming, with boosting being carried out mucosally using a soluble polypeptide or derivative of the invention in combination with a mucosal adjuvant (e.g., LT).

A polypeptide derivative of the invention is also used in accordance with the seventh aspect as a diagnostic
20 reagent for detecting the presence of anti-*Chlamydia* antibodies, e.g., in a blood sample. Such polypeptides are about 5 to about 80, preferably about 10 to about 50 amino acids in length. They are either labeled or unlabeled, depending upon the diagnostic method. Diagnostic methods
25 involving such a reagent are described below.

Upon expression of a DNA molecule of the invention, a polypeptide or polypeptide derivative is produced and purified using known laboratory techniques. As described above, the polypeptide or polypeptide derivative may be produced as a
30 fusion protein containing a fused tail that facilitates purification. The fusion product is used to immunize a small mammal, e.g., a mouse or a rabbit, in order to raise antibodies against the polypeptide or polypeptide derivative (monospecific antibodies). Accordingly, an eighth aspect of the invention

provides a monospecific antibody that binds to a polypeptide or polypeptide derivative of the invention.

By "monospecific antibody" is meant an antibody that is capable of reacting with a unique naturally-occurring
5 *Chlamydia* polypeptide. An antibody of the invention is either polyclonal or monoclonal. Monospecific antibodies may be recombinant, e.g., chimeric (e.g., constituted by a variable region of murine origin associated with a human constant region), humanized (a human immunoglobulin constant backbone
10 together with hypervariable region of animal, e.g., murine, origin), and/or single chain. Both polyclonal and monospecific antibodies may also be in the form of immunoglobulin fragments, e.g., F(ab)'2 or Fab fragments. The antibodies of the invention are of any isotype, e.g., IgG or IgA, and polyclonal
15 antibodies are of a single isotype or a mixture of isotypes.

Antibodies against the polypeptides, homologs or fragments of the present invention are generated by immunization of a mammal with a composition comprising said polypeptide, homolog or fragment. Such antibodies may be
20 polyclonal or monoclonal. Methods to produce polyclonal or monoclonal antibodies are well known in the art. For a review, see "Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, Eds. E. Harlow and D. Lane (1988), and D.E. Yelton et al., 1981. Ann. Rev. Biochem. 50:657-680. For monoclonal
25 antibodies, see Kohler & Milstein (1975) Nature 256:495-497.

The antibodies of the invention, which are raised to a polypeptide or polypeptide derivative of the invention, are produced and identified using standard immunological assays, e.g., Western blot analysis, dot blot assay, or ELISA (see,
30 e.g., Coligan et al., Current Protocols in Immunology (1994) John Wiley & Sons, Inc., New York, NY). The antibodies are used in diagnostic methods to detect the presence of a *Chlamydia* antigen in a sample, such as a biological sample. The antibodies are also used in affinity chromatography for

purifying a polypeptide or polypeptide derivative of the invention. As is discussed further below, such antibodies may be used in prophylactic and therapeutic passive immunization methods.

5 Accordingly, a ninth aspect of the invention provides (i) a reagent for detecting the presence of *Chlamydia* in a biological sample that contains an antibody, polypeptide, or polypeptide derivative of the invention; and (ii) a diagnostic method for detecting the presence of *Chlamydia* in a biological
10 sample, by contacting the biological sample with an antibody, a polypeptide, or a polypeptide derivative of the invention, such that an immune complex is formed, and by detecting such complex to indicate the presence of *Chlamydia* in the sample or the organism from which the sample is derived.

15 Those skilled in the art will readily understand that the immune complex is formed between a component of the sample and the antibody, polypeptide, or polypeptide derivative, whichever is used, and that any unbound material is removed prior to detecting the complex. It is understood that a
20 polypeptide reagent is useful for detecting the presence of anti-*Chlamydia* antibodies in a sample, e.g., a blood sample, while an antibody of the invention is used for screening a sample, such as a gastric extract or biopsy, for the presence of *Chlamydia* polypeptides.

25 For diagnostic applications, the reagent (i.e., the antibody, polypeptide, or polypeptide derivative of the invention) is either in a free state or immobilized on a solid support, such as a tube, a bead, or any other conventional support used in the field. Immobilization is achieved using
30 direct or indirect means. Direct means include passive adsorption (non-covalent binding) or covalent binding between the support and the reagent. By "indirect means" is meant that an anti-reagent compound that interacts with a reagent is first attached to the solid support. For example, if a polypeptide

reagent is used, an antibody that binds to it can serve as an anti-reagent, provided that it binds to an epitope that is not involved in the recognition of antibodies in biological samples. Indirect means may also employ a ligand-receptor system, for example, where a molecule such as a vitamin is grafted onto the polypeptide reagent and the corresponding receptor immobilized on the solid phase. This is illustrated by the biotin-streptavidin system. Alternatively, a peptide tail is added chemically or by genetic engineering to the reagent and the grafted or fused product immobilized by passive adsorption or covalent linkage of the peptide tail.

Such diagnostic agents may be included in a kit which also comprises instructions for use. The reagent is labeled with a detection means which allows for the detection of the reagent when it is bound to its target. The detection means may be a fluorescent agent such as fluorescein isocyanate or fluorescein isothiocyanate, or an enzyme such as horse radish peroxidase or luciferase or alkaline phosphatase, or a radioactive element such as ^{125}I or ^{51}Cr .

Accordingly, a tenth aspect of the invention provides a process for purifying, from a biological sample, a polypeptide or polypeptide derivative of the invention, which involves carrying out antibody-based affinity chromatography with the biological sample, wherein the antibody is a monospecific antibody of the invention.

For use in a purification process of the invention, the antibody is either polyclonal or monospecific, and preferably is of the IgG type. Purified IgGs is prepared from an antiserum using standard methods (see, e.g., Coligan et al., Current Protocols in Immunology (1994) John Wiley & Sons, Inc., New York, NY.). Conventional chromatography supports, as well as standard methods for grafting antibodies, are described in, e.g., Antibodies: A Laboratory Manual, D. Lane, E. Harlow, Eds. (1988) and outlined below.

Briefly, a biological sample, such as an *C. pneumoniae* extract preferably in a buffer solution, is applied to a chromatography material, preferably equilibrated with the buffer used to dilute the biological sample so that the
5 polypeptide or polypeptide derivative of the invention (*i.e.*, the antigen) is allowed to adsorb onto the material. The chromatography material, such as a gel or a resin coupled to an antibody of the invention, is in either a batch form or a column. The unbound components are washed off and the antigen
10 is then eluted with an appropriate elution buffer, such as a glycine buffer or a buffer containing a chaotropic agent, *e.g.*, guanidine HCl, or high salt concentration (*e.g.*, 3 M MgCl₂). Eluted fractions are recovered and the presence of the antigen is detected, *e.g.*, by measuring the absorbance at 280 nm.

15 An eleventh aspect of the invention provides (i) a composition of matter comprising a monospecific antibody of the invention, together with a diluent or carrier; (ii) a pharmaceutical composition comprising a therapeutically or prophylactically effective amount of a monospecific antibody of
20 the invention, and (iii) a method for treating or preventing a *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumoniae* or *C. pecorum*) infection, by administering a therapeutic or prophylactic amount of a monospecific antibody of the invention to an infected individual. Additionally, the eleventh aspect
25 of the invention encompasses the use of a monospecific antibody of the invention in the preparation of a medicament for treating or preventing *Chlamydia* infection.

The monospecific antibody is either polyclonal or monoclonal, preferably of the IgA isotype (predominantly). In
30 passive immunization, the antibody is administered to a mucosal surface of a mammal, *e.g.*, the gastric mucosa, *e.g.*, orally or intragastrically, advantageously, in the presence of a bicarbonate buffer. Alternatively, systemic administration, not requiring a bicarbonate buffer, is carried out. A

monospecific antibody of the invention is administered as a single active component or as a mixture with at least one monospecific antibody specific for a different *Chlamydia* polypeptide. The amount of antibody and the particular regimen used are readily determined by one skilled in the art. For example, daily administration of about 100 to 1,000 mg of antibodies over one week, or three doses per day of about 100 to 1,000 mg of antibodies over two or three days, are effective regimens for most purposes.

Therapeutic or prophylactic efficacy are evaluated using standard methods in the art, e.g., by measuring induction of a mucosal immune response or induction of protective and/or therapeutic immunity, using, e.g., the *C. pneumoniae* mouse model. Those skilled in the art will readily recognize that the *C. pneumoniae* strain of the model may be replaced with another *Chlamydia* strain. For example, the efficacy of DNA molecules and polypeptides from *C. pneumoniae* is preferably evaluated in a mouse model using *C. pneumoniae* strain.

Protection is determined by comparing the degree of *Chlamydia* infection to that of a control group. Protection is shown when infection is reduced by comparison to the control group. Such an evaluation is made for polynucleotides, vaccine vectors, polypeptides and derivatives thereof, as well as antibodies of the invention.

Adjuvants useful in any of the vaccine compositions described above are as follows.

Adjuvants for parenteral administration include aluminum compounds, such as aluminum hydroxide, aluminum phosphate, and aluminum hydroxy phosphate. The antigen is precipitated with, or adsorbed onto, the aluminum compound according to standard protocols. Other adjuvants, such as RIBI (ImmunoChem, Hamilton, MT), are used in parenteral administration.

Adjuvants for mucosal administration include bacterial toxins, e.g., the cholera toxin (CT), the *E. coli* heat-labile toxin (LT), the *Clostridium difficile* toxin A and the pertussis toxin (PT), or combinations, subunits, toxoids, or mutants thereof such as a purified preparation of native cholera toxin subunit B (CTB). Fragments, homologs, derivatives, and fusions to any of these toxins are also suitable, provided that they retain adjuvant activity. Preferably, a mutant having reduced toxicity is used. Suitable mutants are described, e.g., in WO 95/17211 (Arg-7-Lys CT mutant), WO 96/06627 (Arg-192-Gly LT mutant), and WO 95/34323 (Arg-9-Lys and Glu-129-Gly PT mutant). Additional LT mutants that are used in the methods and compositions of the invention include, e.g., Ser-63-Lys, Ala-69Gly, Glu-110-Asp, and Glu-112-Asp mutants. Other adjuvants, such as a bacterial monophosphoryl lipid A (MPLA) of, e.g., *E. coli*, *Salmonella minnesota*, *Salmonella typhimurium*, or *Shigella flexneri*; saponins, or polylactide glycolide (PLGA) microspheres, is also be used in mucosal administration.

Adjuvants useful for both mucosal and parenteral administrations include polyphosphazene (WO 95/02415), DC-chol (3 b-(N-(N',N'-dimethyl aminomethane)-carbamoyl) cholesterol; U.S. Patent No. 5,283,185 and WO 96/14831) and QS-21 (WO 88/09336).

Any pharmaceutical composition of the invention containing a polynucleotide, a polypeptide, a polypeptide derivative, or an antibody of the invention, is manufactured in a conventional manner. In particular, it is formulated with a pharmaceutically acceptable diluent or carrier, e.g., water or a saline solution such as phosphate buffer saline. In general, a diluent or carrier is selected on the basis of the mode and route of administration, and standard pharmaceutical practice. Suitable pharmaceutical carriers or diluents, as well as pharmaceutical necessities for their use in pharmaceutical

formulations, are described in *Remington's Pharmaceutical Sciences*, a standard reference text in this field and in the USP/NF.

The invention also includes methods in which
5 *Chlamydia* infection are treated by oral administration of a
Chlamydia polypeptide of the invention and a mucosal adjuvant,
in combination with an antibiotic, an antacid, sucralfate, or a
combination thereof. Examples of such compounds that can be
administered with the vaccine antigen and the adjuvant are
10 antibiotics, including, e.g., macrolides, tetracyclines, and
derivatives thereof (specific examples of antibiotics that can
be used include azithromycin or doxycycline or immunomodulators
such as cytokines or steroids). In addition, compounds
containing more than one of the above-listed components coupled
15 together, are used. The invention also includes compositions
for carrying out these methods, i.e., compositions containing a
Chlamydia antigen (or antigens) of the invention, an adjuvant,
and one or more of the above-listed compounds, in a
pharmaceutically acceptable carrier or diluent.

20 It has recently been shown that the 60kDa cysteine
rich membrane protein contains a sequence cross-reactive with
the murine alpha-myosin heavy chain epitope M7A-alpha, an
epitope conserved in humans (Bachmaier et al., Science (1999)
283:1335). This cross-reactivity is proposed to contribute to
25 the development of cardiovascular disease, so it may be
beneficial to remove this epitope, and any other epitopes
cross-reactive with human antigens, from the protein if it is
to be used as a vaccine. Accordingly, a further embodiment of
the present invention includes the modification of the coding
30 sequence, for example, by deletion or substitution of the
nucleotides encoding the epitope from polynucleotides encoding
the protein, as to improve the efficacy and safety of the
protein as a vaccine. A similar approach may be appropriate

for any protective antigen found to have unwanted homologies or cross-reactivities with human antigens.

Amounts of the above-listed compounds used in the methods and compositions of the invention are readily
5 determined by one skilled in the art. Treatment/immunization schedules are also known and readily designed by one skilled in the art. For example, the non-vaccine components can be administered on days 1-14, and the vaccine antigen + adjuvant can be administered on days 7, 14, 21, and 28.

10 EXAMPLES

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples. These examples are described solely for purposes of illustration and are not
15 intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of
20 limitation.

Example 1:

This example illustrates the preparation of a plasmid vector pCABk1043 containing the lpxB gene.

The lpxB gene was amplified from *Chlamydia pneumoniae*
25 genomic DNA by polymerase chain reaction (PCR) using a 5' primer (5' ATAAGAATGCGGCCGCCACCATGATCCCTTCTGGCCTAGTTTATC 3'; SEQ ID No:3 and a 3' primer (5' CGGGGTACCGAATATTCATACAGCGGGAAGTGTG 3'; SEQ ID No:4). The
30 5' primer contains a Not I restriction site, a ribosome binding site, an initiation codon and a sequence at the 5' end of the lpxB coding sequence. The 3' primer includes the sequence encoding the C-terminal sequence of the lpxB and a Kpn I restriction site. The stop codon was excluded and an additional

nucleotide was inserted to obtain an in-frame fusion with the Histidine tag.

After amplification, the PCR fragment was purified using QIAquick™ PCR purification kit (Qiagen) and then digested with Not I and Kpn I and cloned into the pCA-Myc-His eukaryotic expression vector describe in Example 2 (Figure 3) with transcription under control of the human CMV promoter.

Example 2:

This example illustrates the preparation of the eukaryotic expression vector pCA/Myc-His.

Plasmid pcDNA3.1(-)Myc-His C (Invitrogen) was restricted with Spe I and Bam HI to remove the CMV promoter and the remaining vector fragment was isolated. The CMV promoter and intron A from plasmid VR-1012 (Vical) was isolated on a Spe I / Bam HI fragment. The fragments were ligated together to produce plasmid pCA/Myc-His. The Not I/Kpn I restricted PCR fragment containing the lpxB gene was ligated into the Not I and Kpn I restricted plasmid pCA/Myc-His to produce plasmid pCABk1043 (Figure 3).

The resulting plasmid, pCABk1043, was transferred by electroporation into *E. coli* XL-1 blue (Stratagene) which was grown in LB broth containing 50 µg/ml of carbenicillin. The plasmid was isolated by Endo Free Plasmid Giga Kit™ (Qiagen) large scale DNA purification system. DNA concentration was determined by absorbance at 260 nm and the plasmid was verified after gel electrophoresis and Ethidium bromide staining and comparison to molecular weight standards. The 5' and 3' ends of the gene were verified by sequencing using a LiCor model 4000 L DNA sequencer and IRD-800 labelled primers.

Example 3:

This example illustrates the immunization of mice to achieve protection against an intranasal challenge of *C. pneumoniae*.

5 It has been previously demonstrated (Yang et. al., 1993) that mice are susceptible to intranasal infection with different isolates of *C. pneumoniae*. Strain AR-39 (Grayston, 1989) was used in Balb/c mice as a challenge infection model to examine the capacity of chlamydia gene products delivered as
10 naked DNA to elicit a protective response against a sublethal *C. pneumoniae* lung infection. Protective immunity is defined as an accelerated clearance of pulmonary infection.

Groups of 7 to 9 week old male Balb/c mice (6 to 10 per group) were immunized intramuscularly (i.m.) plus
15 intranasally (i.n.) with plasmid DNA containing the coding sequence of *C. pneumoniae* lpxB as described in Examples 1 and 2. Saline or the plasmid vector lacking an inserted chlamydial gene was given to groups of control animals.

For i.m. immunization, alternate left and right
20 quadriceps were injected with 100µg of DNA in 50µl of PBS on three occasions at 0, 3 and 6 weeks. For i.n. immunization, anaesthetized mice aspirated 50µl of PBS containing 50 µg DNA on three occasions at 0, 3 and 6 weeks. At week 8, immunized mice were inoculated i.n. with 5×10^5 IFU of *C. pneumoniae*,
25 strain AR39 in 100µl of SPG buffer to test their ability to limit the growth of a sublethal *C. pneumoniae* challenge.

Lungs were taken from mice at day 9 post-challenge and immediately homogenised in SPG buffer (7.5% sucrose, 5mM glutamate, 12.5mM phosphate pH 7.5). The homogenate was stored
30 frozen at -70°C until assay. Dilutions of the homogenate were assayed for the presence of infectious chlamydia by inoculation onto monolayers of susceptible cells. The inoculum was centrifuged onto the cells at 3000rpm for 1 hour, then the

cells were incubated for three days at 35°C in the presence of 1µg/ml cycloheximide. After incubation the monolayers were fixed with formalin and methanol then immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with *C. pneumoniae* and metal-enhanced DAB as a peroxidase substrate.

Figure 4 and Table 1 show that mice immunized i.n. and i.m. with pCABk1043 had chlamydial lung titers less than 37,000 in 5 of 6 cases at day 9 (mean 39,666) whereas the range of values for control mice sham immunized with saline was 13,600-458,100 IFU/lung (mean 107,641) at day 9. DNA immunisation *per se* was not responsible for the observed protective effect since another plasmid DNA construct, pCABk917, failed to protect, with lung titers in immunised mice similar (mean 85,350 IFU/lung) to those obtained for saline-immunized control mice. The construct pCABk917 is identical to pCABk1043 except that the nucleotide sequence encoding the lpxB is replaced with a *C. pneumoniae* nucleotide sequence encoding an outer membrane protein.

Table 1

MOUSE	BACTERIAL LOAD (INCLUSION FORMING UNITS PER LUNG) IN THE LUNGS OF BALB/C MICE IMMUNIZED WITH VARIOUS DNA IMMUNIZATION CONSTRUCTS		
	IMMUNIZING CONSTRUCT		
	Saline	pCABk917	pCABk1043
	Day 9	Day 9	Day 9
1	90000	72400	26600
2	69600	108000	27500
3	136400	24700	26000
4	458100	0	32500
5	166500	83600	89300
6	49500	223400	36100
7	13600		
8	150600		
9	37600		
10	179700		
11	91100		
12	289400		
13	16200		
14	233300		
15	36300		
16	132700		
17	57300		
18	36900		
19	115000		
20	108000		
21	32600		
22	33400		
23	16300		
24	33300		
MEAN	107641.7	85350	39666.67
SD	104011.6	78364.07	24629.63
Wilcoxon p		0.6223	0.03142

CLAIMS:

1. A nucleic acid molecule comprising a nucleic acid sequence which encodes a polypeptide selected from any of:
 - (a) SEQ ID No: 2;
 - 5 (b) an immunogenic fragment comprising at least 12 consecutive amino acids from a polypeptide of (a); and
 - (c) a polypeptide of (a) or (b) which has been modified to improve its immunogenicity, wherein said modified polypeptide is at least 75% identical in amino acid sequence to
 - 10 the corresponding polypeptide of (a) or (b).
2. A nucleic acid molecule comprising a nucleic acid sequence selected from any of:
 - (a) SEQ ID No: 1;
 - (b) a sequence which encodes a polypeptide encoded
 - 15 by SEQ ID No: 1;
 - (c) a sequence comprising at least 38 consecutive nucleotides from any one of the nucleic acid sequences of (a) and (b); and
 - (d) a sequence which encodes a polypeptide which is
 - 20 at least 75% identical in amino acid sequence to the polypeptides encoded by SEQ ID No: 1.
3. A nucleic acid molecule comprising a nucleic acid sequence which is anti-sense to the nucleic acid molecule of claim 1.
- 25 4. A nucleic acid molecule comprising a nucleic acid sequence which encodes a fusion protein, said fusion protein comprising a polypeptide encoded by a nucleic acid molecule according to claim 1 and an additional polypeptide.
5. The nucleic acid molecule of claim 4 wherein the
- 30 additional polypeptide is a heterologous signal peptide.

6. The nucleic acid molecule of claim 4 wherein the additional polypeptide has adjuvant activity.

7. A nucleic acid molecule according to any one of claims 1 to 6, operatively linked to one or more expression
5 control sequences.

8. A vaccine comprising at least one first nucleic acid according to any one of claims 1, 2, and 4 to 7 and a vaccine vector wherein each first nucleic acid is expressed as a polypeptide, the vaccine optionally comprising a second nucleic
10 acid encoding an additional polypeptide which enhances the immune response to the polypeptide expressed by said first nucleic acid.

9. The vaccine of claim 8 wherein the second nucleic acid encodes an additional *Chlamydia* polypeptide.

15 10. A pharmaceutical composition comprising a nucleic acid according to any one of claims 1 to 7 and a pharmaceutically acceptable carrier.

11. A pharmaceutical composition comprising a vaccine according to claim 8 or 9 and a pharmaceutically acceptable
20 carrier.

12. A unicellular host transformed with the nucleic acid molecule of claim 7.

13. A nucleic acid probe of 5 to 100 nucleotides which hybridizes under stringent conditions to the nucleic acid
25 molecule of SEQ ID No: 1, or to a homolog or complementary or anti-sense sequence of said nucleic acid molecule.

14. A primer of 10 to 40 nucleotides which hybridizes under stringent conditions to the nucleic acid molecules of SEQ

ID No: 1, or to a homolog or complementary or anti-sense sequence of said nucleic acid molecule.

15. A polypeptide encoded by a nucleic acid sequence according to any one of claims 1, 2 and 4 to 7.

5 16. A polypeptide comprising an amino acid sequence selected from any of:

(a) SEQ ID No: 2;

(b) an immunogenic fragment comprising at least 12 consecutive amino acids from a polypeptide of (a); and

10 (c) a polypeptide of (a) or (b) which has been modified to improve its immunogenicity, wherein said modified polypeptide is at least 75% identical in amino acid sequence to the corresponding polypeptide of (a) or (b).

15 17. A fusion polypeptide comprising a polypeptide of claim 15 or 16 and an additional polypeptide.

18. The fusion polypeptide of claim 17 wherein the additional polypeptide is a heterologous signal peptide.

19. The fusion protein of claim 17 wherein the additional polypeptide has adjuvant activity.

20 20. A method for producing a polypeptide of claim 15 or 16, comprising the step of culturing a unicellular host according to claim 12.

21. An antibody against the polypeptide of any one of claims 15 to 19.

25 22. A vaccine comprising at least one first polypeptide according to any one of claims 15 to 19 and a pharmaceutically acceptable carrier, optionally comprising a second polypeptide which enhances the immune response to the first polypeptide.

23. The vaccine of claim 22 wherein the second polypeptide comprises an additional *Chlamydia* polypeptide.

24. A pharmaceutical composition comprising a polypeptide according to any one of claims 15 to 19 and a pharmaceutically acceptable carrier.

25. A pharmaceutical composition comprising a vaccine according to claim 22 or 23 and a pharmaceutically acceptable carrier.

26. A pharmaceutical composition comprising an antibody according to claim 21 and a pharmaceutically acceptable carrier.

27. A method for preventing or treating *Chlamydia* infection using:

- (a) the nucleic acid of any one of claims 1 to 7;
- 15 (b) the vaccine of any one of claims 8, 9, 22 and 23;
- (c) the pharmaceutical composition of any one of claims 10, 11, 24 to 26;
- (d) the polypeptide of any one of claims 15 to 19;
- 20 or
- (e) the antibody of claim 21.

28. A method of detecting *Chlamydia* infection comprising the step of assaying a body fluid of a mammal to be tested, with a component selected from any one of:

- 25 (a) the nucleic acid of any one of claims 1 to 7;
- (b) the polypeptide of any one of claims 15 to 19;
- and
- (c) the antibody of claim 21.

29. A diagnostic kit comprising instructions for use and a component selected from any one of:

- (a) the nucleic acid of any one of claims 1 to 7;
- (b) the polypeptide of any one of claims 15 to 19;

and

- (c) the antibody of claim 21.

- 5 30. A method for identifying a polypeptide of claims 15 to 19 which induces an immune response effective to prevent or lessen the severity of *Chlamydia* infection in a mammal previously immunized with polypeptide, comprising the steps of:
- (a) immunizing a mouse with the polypeptide; and
 - 10 (b) inoculating the immunized mouse with *Chlamydia*;
- wherein the polypeptide which prevents or lessens the severity of *Chlamydia* infection in the immunized mouse compared to a non-immunized control mouse is identified.

31. Expression plasmid pCABk1043.

15 32. A nucleic acid molecule of SEQ ID NO. 3 or 4.

33. lpxB from *Chlamydia*.

34. lpxB from *Chlamydia pneumoniae*.

1/12

Figure 1. Sequence of *C. pneumoniae* lpxB gene.

5	tagtgtatcc taagctcgat aaagtgtatg ttgcttggat acgtcatcat caaacactta	60
	agtgtaaaaa ggattctcat tctcaaaagt aggtctaaac atg atc cct tct ggc	115
		Met Ile Pro Ser Gly 1 5
10	cta gtt tat ctg ctt tat ccc cta gga ttt ctt gct agt ctt ttt ttt	163
	Leu Val Tyr Leu Leu Tyr Pro Leu Gly Phe Leu Ala Ser Leu Phe Phe	10 15 20
15	ggg agt gcc ttc tca ata cag tgg tgg ctg agt aaa aag cgc aaa gag	211
	Gly Ser Ala Phe Ser Ile Gln Trp Trp Leu Ser Lys Lys Arg Lys Glu	25 30 35
20	gtc tat gct cct cga agc ttt tgg att ctc tct tca ata gga gcc act	259
	Val Tyr Ala Pro Arg Ser Phe Trp Ile Leu Ser Ser Ile Gly Ala Thr	40 45 50
25	ttg atg att gtc cac ggg aca att caa agc caa ttt cct gtg act gta	307
	Leu Met Ile Val His Gly Thr Ile Gln Ser Gln Phe Pro Val Thr Val	55 60 65
30	ctt cat gtc att aac tta att atc tac ctc aga aac cta aat att act	355
	Leu His Val Ile Asn Leu Ile Ile Tyr Leu Arg Asn Leu Asn Ile Thr	70 75 80 85
35	tcc tct cga ccg att tcc ttt cga gcg acc tta gtt ctg atg gct ctg	403
	Ser Ser Arg Pro Ile Ser Phe Arg Ala Thr Leu Val Leu Met Ala Leu	90 95 100
40	atg gcc tct cct aat atc ttc cat ctt cct ctt cct cct gca caa ctt	499
	Met Ala Ser Pro Asn Ile Phe His Leu Pro Leu Pro Pro Ala Gln Leu	120 125 130
45	tct tgg cat ctt ata ggc tgt tta ggg ctt gct att ttt tcg ggt cgt	547
	Ser Trp His Leu Ile Gly Cys Leu Gly Leu Ala Ile Phe Ser Gly Arg	135 140 145
50	ttt tta atc caa tgg ttc tac ata gaa tcg aat aat act aaa gat ttc	595
	Phe Leu Ile Gln Trp Phe Tyr Ile Glu Ser Asn Asn Thr Lys Asp Phe	150 155 160 165
55	ccc tta cta ttt tgg aaa att gga ctt ctt ggg gga ttg ctg gcg ctg	643
	Pro Leu Leu Phe Trp Lys Ile Gly Leu Leu Gly Gly Leu Leu Ala Leu	170 175 180
60	gtg tac ttc att cgc atc gga gat cct ata aac atc ctg tgc tat ggc	691
	Val Tyr Phe Ile Arg Ile Gly Asp Pro Ile Asn Ile Leu Cys Tyr Gly	185 190 195
	tgt gga tta ttc ccg tcc atc gca aat tta cgt ctc ttt tat aaa gaa	739
	Cys Gly Leu Phe Pro Ser Ile Ala Asn Leu Arg Leu Phe Tyr Lys Glu	200 205 210

2/12

Figure 1 (Continued)


5	caa cgt tcg act cct tat ttg gat aca cac tgc ttt ctc tct gct gga 787 Gln Arg Ser Thr Pro Tyr Leu Asp Thr His Cys Phe Leu Ser Ala Gly 215 220 225
10	gaa gct agc ggt gac att ttg ggg ggt aag tta atc caa tcc ata aag 835 Glu Ala Ser Gly Asp Ile Leu Gly Gly Lys Leu Ile Gln Ser Ile Lys 230 235 240 245
15	tct ctc tat cca aac ata cgg ttc tgg ggt gtt gga ggt cct gca atg 883 Ser Leu Tyr Pro Asn Ile Arg Phe Trp Gly Val Gly Gly Pro Ala Met 250 255 260
20	cga caa gaa ggt ctc cag ccc att cta aac atg gag gag ttt caa gtt 931 Arg Gln Glu Gly Leu Gln Pro Ile Leu Asn Met Glu Glu Phe Gln Val 265 270 275
25	tca gga ttt gct gaa gtt ctg ggg tca tta ttt aga tta tat cga aat 979 Ser Gly Phe Ala Glu Val Leu Gly Ser Leu Phe Arg Leu Tyr Arg Asn 280 285 290
30	tac cgc aaa atc cta aaa acg ata ctc aaa cac aag cca gcg act ctt 1027 Tyr Arg Lys Ile Leu Lys Thr Ile Leu Lys His Lys Pro Ala Thr Leu 295 300 305
35	atc ttc att gat ttc ccc gat ttt cac ctt ctc tta att aaa aag ctg 1075 Ile Phe Ile Asp Phe Pro Asp Phe His Leu Leu Leu Ile Lys Lys Leu 310 315 320 325
40	aga aaa cat ggg tat cga gga aaa atc atc cat tac gtt tgc ccg agt 1123 Arg Lys His Gly Tyr Arg Gly Lys Ile Ile His Tyr Val Cys Pro Ser 330 335 340
45	att tgg gca tgg cgc ccc aaa aga aaa cgc ata cta gag caa cat ctt 1171 Ile Trp Ala Trp Arg Pro Lys Arg Lys Arg Ile Leu Glu Gln His Leu 345 350 355
50	gat atg ctc cta ctg atc ctt cct ttt gaa gaa ggc ctt ttc aaa aac 1219 Asp Met Leu Leu Leu Ile Leu Pro Phe Glu Glu Gly Leu Phe Lys Asn 360 365 370
55	aca tct ctt gag act gtg tat tta gga cac ccc cta gtt gag gag atc 1267 Thr Ser Leu Glu Thr Val Tyr Leu Gly His Pro Leu Val Glu Glu Ile 375 380 385
60	tct gac tac aaa gaa cag gcc tct tgg aaa gag aaa ttc ctt aat tct 1315 Ser Asp Tyr Lys Glu Gln Ala Ser Trp Lys Glu Lys Phe Leu Asn Ser 390 395 400 405
65	gac cgt cct att gtt gct gcg ttt cct gga agt cgt cgt ggt gac atc 1363 Asp Arg Pro Ile Val Ala Ala Phe Pro Gly Ser Arg Arg Gly Asp Ile 410 415 420
70	agc cga aat tta cgg atc cag gta caa gct ttt cta aat tca tcg cta 1411 Ser Arg Asn Leu Arg Ile Gln Val Gln Ala Phe Leu Asn Ser Ser Leu 425 430 435
75	tct caa aca cat caa ttt gtg gtt tcc tca tca agc gct aag tac gac 1459 Ser Gln Thr His Gln Phe Val Val Ser Ser Ser Ser Ala Lys Tyr Asp 440 445 450

3/12

Figure 1 (Continued)


5	gaa atc att gaa gat act cta aaa gca gaa gga tgt cag cat agt caa 1507 Glu Ile Ile Glu Asp Thr Leu Lys Ala Glu Gly Cys Gln His Ser Gln 455 460 465
10	atc atc ccc atg aat ttc cgc tat gag ctc atg aga agc tgt gat tgt 1555 Ile Ile Pro Met Asn Phe Arg Tyr Glu Leu Met Arg Ser Cys Asp Cys 470 475 480 485
15	gca cta gca aaa tgt ggc act ata gtc tta gaa act gcc tta aat caa 1603 Ala Leu Ala Lys Cys Gly Thr Ile Val Leu Glu Thr Ala Leu Asn Gln 490 495 500
20	aca ccc aca ata gtc atg tgt cga ctt cga cct ttt gac aca ttc tta 1651 Thr Pro Thr Ile Val Met Cys Arg Leu Arg Pro Phe Asp Thr Phe Leu 505 510 515
25	gca aaa tat att ttt aaa att ctt ttg ccc gcg tat tcc tta cca aat 1699 Ala Lys Tyr Ile Phe Lys Ile Leu Leu Pro Ala Tyr Ser Leu Pro Asn 520 525 530
30	atc att atg aac tcc gtg att ttt cca gag ttc att gga ggg aaa aaa 1747 Ile Ile Met Asn Ser Val Ile Phe Pro Glu Phe Ile Gly Gly Lys Lys 535 540 545
35	gat ttt cat cca gaa gaa atc gct acc gct tta gat ttg cta aat caa 1795 Asp Phe His Pro Glu Glu Ile Ala Thr Ala Leu Asp Leu Leu Asn Gln 550 555 560 565
40	cat ggc tct aaa gaa aaa caa aag gag gac tgc cgc aaa tta tgt aaa 1843 His Gly Ser Lys Glu Lys Gln Lys Glu Asp Cys Arg Lys Leu Cys Lys 570 575 580
45	gtg atg acg aca ggg caa ata gct tct gaa gag ttt tta aaa aga att 1891 Val Met Thr Thr Gly Gln Ile Ala Ser Glu Glu Phe Leu Lys Arg Ile 585 590 595
50	ttt gac aca ctt ccc gct gta tga ata ttt aggtttctta aaacaagtgc 1941 Phe Asp Thr Leu Pro Ala Val 600 605
	tttaagaact cataaataag tataaaacaa ttcgttttga gctacttttt tatagttatt 2001
	ctaaataaaaa attaatgaa ta 2023

4/12

Figure 2. Restriction enzyme analysis of the *C. pneumoniae* lpxB gene.


5/12

Figure 2 (Continued)




6/12

Figure 2 (Continued)




7/12

Figure 2 (Continued)



8/12

Figure 2 (Continued)



9/12

Figure 2 (Continued)





Figure 2 (Continued)







FIG. 3
Construction of pCABk1043

12/12

Figure 4. Protective efficacy of DNA immunization with pCABk1043



1/6
SEQUENCE LISTING

<110> Aventis Pasteur Limited

5 <120> Chlamydia antigens and corresponding DNA fragments and uses thereof

<130> 77813-22

<140>

10 <141>

<150> US 60/154,461

<151> 1999-09-17

15 <160> 4

<170> PatentIn Ver. 2.0

<210> 1

20 <211> 2023

<212> DNA

<213> Chlamydia pneumoniae

<220>

25 <221> CDS

<222> (101)..(1921)

<400> 1

30 tagtgtatcc taagctcgat aaagtgtatg ttgcttggat acgtcatcat caaacactta 60

agtgtaaaaa ggattctcat tctcaaaagt aggtctaaac atg atc cct tct ggc 115

Met Ile Pro Ser Gly

1 5

35 cta gtt tat ctg ctt tat ccc cta gga ttt ctt gct agt ctt ttt ttt 163

Leu Val Tyr Leu Leu Tyr Pro Leu Gly Phe Leu Ala Ser Leu Phe Phe

10 15 20

40 ggg agt gcc ttc tca ata cag tgg tgg ctg agt aaa aag cgc aaa gag 211

Gly Ser Ala Phe Ser Ile Gln Trp Trp Leu Ser Lys Lys Arg Lys Glu

25 30 35

45 gtc tat gct cct cga agc ttt tgg att ctc tct tca ata gga gcc act 259

Val Tyr Ala Pro Arg Ser Phe Trp Ile Leu Ser Ser Ile Gly Ala Thr

40 45 50

50 ttg atg att gtc cac ggg aca att caa agc caa ttt cct gtg act gta 307

Leu Met Ile Val His Gly Thr Ile Gln Ser Gln Phe Pro Val Thr Val

55 60 65

55 ctt cat gtc att aac tta att atc tac ctc aga aac cta aat att act 355

Leu His Val Ile Asn Leu Ile Ile Tyr Leu Arg Asn Leu Asn Ile Thr

70 75 80 85

60 tcc tct cga ccg att tcc ttt cga gcg acc tta gtt ctg atg gct ctg 403

Ser Ser Arg Pro Ile Ser Phe Arg Ala Thr Leu Val Leu Met Ala Leu

90 95 100

60 agc gtg gtg ttc gtg act ctg cct ttc ctc tat gtg aac atg gaa tgg 451

Ser Val Val Phe Val Thr Leu Pro Phe Leu Tyr Val Asn Met Glu Trp

105 110 115

2/6

	atg gcc tct cct aat atc ttc cat ctt cct ctt cct cct gca caa ctt	499
	Met Ala Ser Pro Asn Ile Phe His Leu Pro Leu Pro Pro Ala Gln Leu	
	120 125 130	
5	tct tgg cat ctt ata ggc tgt tta ggg ctt gct att ttt tcg ggt cgt	547
	Ser Trp His Leu Ile Gly Cys Leu Gly Leu Ala Ile Phe Ser Gly Arg	
	135 140 145	
10	ttt tta atc caa tgg ttc tac ata gaa tcg aat aat act aaa gat ttc	595
	Phe Leu Ile Gln Trp Phe Tyr Ile Glu Ser Asn Asn Thr Lys Asp Phe	
	150 155 160	
15	ccc tta cta ttt tgg aaa att gga ctt ctt ggg gga ttg ctg gcg ctg	643
	Pro Leu Leu Phe Trp Lys Ile Gly Leu Leu Gly Gly Leu Leu Ala Leu	
	170 175 180	
20	gtg tac ttc att cgc atc gga gat cct ata aac atc ctg tgc tat ggc	691
	Val Tyr Phe Ile Arg Ile Gly Asp Pro Ile Asn Ile Leu Cys Tyr Gly	
	185 190 195	
25	tgt gga tta ttc ccg tcc atc gca aat tta cgt ctc ttt tat aaa gaa	739
	Cys Gly Leu Phe Pro Ser Ile Ala Asn Leu Arg Leu Phe Tyr Lys Glu	
	200 205 210	
30	caa cgt tcg act cct tat ttg gat aca cac tgc ttt ctc tct gct gga	787
	Gln Arg Ser Thr Pro Tyr Leu Asp Thr His Cys Phe Leu Ser Ala Gly	
	215 220 225	
35	gaa gct agc ggt gac att ttg ggg ggt aag tta atc caa tcc ata aag	835
	Glu Ala Ser Gly Asp Ile Leu Gly Gly Lys Leu Ile Gln Ser Ile Lys	
	230 235 240 245	
40	tct ctc tat cca aac ata cgg ttc tgg ggt gtt gga ggt cct gca atg	883
	Ser Leu Tyr Pro Asn Ile Arg Phe Trp Gly Val Gly Gly Pro Ala Met	
	250 255 260	
45	cga caa gaa ggt ctc cag ccc att cta aac atg gag gag ttt caa gtt	931
	Arg Gln Glu Gly Leu Gln Pro Ile Leu Asn Met Glu Glu Phe Gln Val	
	265 270 275	
50	tca gga ttt gct gaa gtt ctg ggg tca tta ttt aga tta tat cga aat	979
	Ser Gly Phe Ala Glu Val Leu Gly Ser Leu Phe Arg Leu Tyr Arg Asn	
	280 285 290	
55	tac cgc aaa atc cta aaa acg ata ctc aaa cac aag cca gcg act ctt	1027
	Tyr Arg Lys Ile Leu Lys Thr Ile Leu Lys His Lys Pro Ala Thr Leu	
	295 300 305	
60	atc ttc att gat ttc ccc gat ttt cac ctt ctc tta att aaa aag ctg	1075
	Ile Phe Ile Asp Phe Pro Asp Phe His Leu Leu Leu Ile Lys Lys Leu	
	310 315 320 325	
65	aga aaa cat ggg tat cga gga aaa atc atc cat tac gtt tgc ccg agt	1123
	Arg Lys His Gly Tyr Arg Gly Lys Ile Ile His Tyr Val Cys Pro Ser	
	330 335 340	
70	att tgg gca tgg cgc ccc aaa aga aaa cgc ata cta gag caa cat ctt	1171
	Ile Trp Ala Trp Arg Pro Lys Arg Lys Arg Ile Leu Glu Gln His Leu	
	345 350 355	

3/6

5	gat atg ctc cta ctg atc ctt cct ttt gaa gaa ggc ctt ttc aaa aac 1219 Asp Met Leu Leu Ile Leu Pro Phe Glu Glu Gly Leu Phe Lys Asn 360 365 370
10	aca tct ctt gag act gtg tat tta gga cac ccc cta gtt gag gag atc 1267 Thr Ser Leu Glu Thr Val Tyr Leu Gly His Pro Leu Val Glu Glu Ile 375 380 385
15	tct gac tac aaa gaa cag gcc tct tgg aaa gag aaa ttc ctt aat tct 1315 Ser Asp Tyr Lys Glu Gln Ala Ser Trp Lys Glu Lys Phe Leu Asn Ser 390 395 400 405
20	gac cgt cct att gtt gct gcg ttt cct gga agt cgt cgt ggt gac atc 1363 Asp Arg Pro Ile Val Ala Ala Phe Pro Gly Ser Arg Arg Gly Asp Ile 410 415 420
25	agc cga aat tta cgg atc cag gta caa gct ttt cta aat tca tcg cta 1411 Ser Arg Asn Leu Arg Ile Gln Val Gln Ala Phe Leu Asn Ser Ser Leu 425 430 435
30	tct caa aca cat caa ttt gtg gtt tcc tca tca agc gct aag tac gac 1459 Ser Gln Thr His Gln Phe Val Val Ser Ser Ser Ser Ala Lys Tyr Asp 440 445 450
35	gaa atc att gaa gat act cta aaa gca gaa gga tgt cag cat agt caa 1507 Glu Ile Ile Glu Asp Thr Leu Lys Ala Glu Gly Cys Gln His Ser Gln 455 460 465
40	atc atc ccc atg aat ttc cgc tat gag ctc atg aga agc tgt gat tgt 1555 Ile Ile Pro Met Asn Phe Arg Tyr Glu Leu Met Arg Ser Cys Asp Cys 470 475 480 485
45	gca cta gca aaa tgt ggc act ata gtc tta gaa act gcc tta aat caa 1603 Ala Leu Ala Lys Cys Gly Thr Ile Val Leu Glu Thr Ala Leu Asn Gln 490 495 500
50	aca ccc aca ata gtc atg tgt cga ctt cga cct ttt gac aca ttc tta 1651 Thr Pro Thr Ile Val Met Cys Arg Leu Arg Pro Phe Asp Thr Phe Leu 505 510 515
55	gca aaa tat att ttt aaa att ctt ttg ccc gcg tat tcc tta cca aat 1699 Ala Lys Tyr Ile Phe Lys Ile Leu Leu Pro Ala Tyr Ser Leu Pro Asn 520 525 530
60	atc att atg aac tcc gtg att ttt cca gag ttc att gga ggg aaa aaa 1747 Ile Ile Met Asn Ser Val Ile Phe Pro Glu Phe Ile Gly Gly Lys Lys 535 540 545
65	gat ttt cat cca gaa gaa atc gct acc gct tta gat ttg cta aat caa 1795 Asp Phe His Pro Glu Glu Ile Ala Thr Ala Leu Asp Leu Leu Asn Gln 550 555 560 565
70	cat ggc tct aaa gaa aaa caa aag gag gac tgc cgc aaa tta tgt aaa 1843 His Gly Ser Lys Glu Lys Gln Lys Glu Asp Cys Arg Lys Leu Cys Lys 570 575 580
75	gtg atg acg aca ggg caa ata gct tct gaa gag ttt tta aaa aga att 1891 Val Met Thr Thr Gly Gln Ile Ala Ser Glu Glu Phe Leu Lys Arg Ile 585 590 595

4/6

```

    ttt gac aca ctt ccc gct gta tga ata ttt aggtttctta aaacaagttc      1941
    Phe Asp Thr Leu Pro Ala Val      Ile Phe
        600                      605

5   ttttaagaact cataaataag tataaaacaa ttcgttttga gctacttttt tatagtattt 2001

    ctaaataaaaa attaattgaa ta                                          2023

10  <210> 2
    <211> 604
    <212> PRT
    <213> Chlamydia pneumoniae

15  <400> 2
    Met Ile Pro Ser Gly Leu Val Tyr Leu Leu Tyr Pro Leu Gly Phe Leu
        1           5           10           15

20  Ala Ser Leu Phe Phe Gly Ser Ala Phe Ser Ile Gln Trp Trp Leu Ser
        20           25           30

    Lys Lys Arg Lys Glu Val Tyr Ala Pro Arg Ser Phe Trp Ile Leu Ser
        35           40           45

25  Ser Ile Gly Ala Thr Leu Met Ile Val His Gly Thr Ile Gln Ser Gln
        50           55           60

    Phe Pro Val Thr Val Leu His Val Ile Asn Leu Ile Ile Tyr Leu Arg
    30   65           70           75           80

    Asn Leu Asn Ile Thr Ser Ser Arg Pro Ile Ser Phe Arg Ala Thr Leu
        85           90           95

35  Val Leu Met Ala Leu Ser Val Val Phe Val Thr Leu Pro Phe Leu Tyr
        100          105          110

    Val Asn Met Glu Trp Met Ala Ser Pro Asn Ile Phe His Leu Pro Leu
        115          120          125

40  Pro Pro Ala Gln Leu Ser Trp His Leu Ile Gly Cys Leu Gly Leu Ala
        130          135          140

    Ile Phe Ser Gly Arg Phe Leu Ile Gln Trp Phe Tyr Ile Glu Ser Asn
    45  145          150          155          160

    Asn Thr Lys Asp Phe Pro Leu Leu Phe Trp Lys Ile Gly Leu Leu Gly
        165          170          175

50  Gly Leu Leu Ala Leu Val Tyr Phe Ile Arg Ile Gly Asp Pro Ile Asn
        180          185          190

    Ile Leu Cys Tyr Gly Cys Gly Leu Phe Pro Ser Ile Ala Asn Leu Arg
        195          200          205

55  Leu Phe Tyr Lys Glu Gln Arg Ser Thr Pro Tyr Leu Asp Thr His Cys
        210          215          220

    Phe Leu Ser Ala Gly Glu Ala Ser Gly Asp Ile Leu Gly Gly Lys Leu
    60  225          230          235          240

```

5/6

	Ile	Gln	Ser	Ile	Lys	Ser	Leu	Tyr	Pro	Asn	Ile	Arg	Phe	Trp	Gly	Val	
					245					250					255		
5	Gly	Gly	Pro	Ala	Met	Arg	Gln	Glu	Gly	Leu	Gln	Pro	Ile	Leu	Asn	Met	
				260					265					270			
	Glu	Glu	Phe	Gln	Val	Ser	Gly	Phe	Ala	Glu	Val	Leu	Gly	Ser	Leu	Phe	
			275					280					285				
10	Arg	Leu	Tyr	Arg	Asn	Tyr	Arg	Lys	Ile	Leu	Lys	Thr	Ile	Leu	Lys	His	
		290					295					300					
	Lys	Pro	Ala	Thr	Leu	Ile	Phe	Ile	Asp	Phe	Pro	Asp	Phe	His	Leu	Leu	
15		305				310					315					320	
	Leu	Ile	Lys	Lys	Leu	Arg	Lys	His	Gly	Tyr	Arg	Gly	Lys	Ile	Ile	His	
					325					330					335		
20	Tyr	Val	Cys	Pro	Ser	Ile	Trp	Ala	Trp	Arg	Pro	Lys	Arg	Lys	Arg	Ile	
				340					345					350			
	Leu	Glu	Gln	His	Leu	Asp	Met	Leu	Leu	Leu	Ile	Leu	Pro	Phe	Glu	Glu	
			355					360					365				
25	Gly	Leu	Phe	Lys	Asn	Thr	Ser	Leu	Glu	Thr	Val	Tyr	Leu	Gly	His	Pro	
		370					375					380					
	Leu	Val	Glu	Glu	Ile	Ser	Asp	Tyr	Lys	Glu	Gln	Ala	Ser	Trp	Lys	Glu	
30		385				390					395					400	
	Lys	Phe	Leu	Asn	Ser	Asp	Arg	Pro	Ile	Val	Ala	Ala	Phe	Pro	Gly	Ser	
				405						410					415		
35	Arg	Arg	Gly	Asp	Ile	Ser	Arg	Asn	Leu	Arg	Ile	Gln	Val	Gln	Ala	Phe	
				420					425					430			
	Leu	Asn	Ser	Ser	Leu	Ser	Gln	Thr	His	Gln	Phe	Val	Val	Ser	Ser	Ser	
		435						440					445				
40	Ser	Ala	Lys	Tyr	Asp	Glu	Ile	Ile	Glu	Asp	Thr	Leu	Lys	Ala	Glu	Gly	
		450					455					460					
	Cys	Gln	His	Ser	Gln	Ile	Ile	Pro	Met	Asn	Phe	Arg	Tyr	Glu	Leu	Met	
45		465				470					475					480	
	Arg	Ser	Cys	Asp	Cys	Ala	Leu	Ala	Lys	Cys	Gly	Thr	Ile	Val	Leu	Glu	
				485						490					495		
50	Thr	Ala	Leu	Asn	Gln	Thr	Pro	Thr	Ile	Val	Met	Cys	Arg	Leu	Arg	Pro	
				500					505					510			
	Phe	Asp	Thr	Phe	Leu	Ala	Lys	Tyr	Ile	Phe	Lys	Ile	Leu	Leu	Pro	Ala	
			515					520					525				
55	Tyr	Ser	Leu	Pro	Asn	Ile	Ile	Met	Asn	Ser	Val	Ile	Phe	Pro	Glu	Phe	
		530					535					540					
	Ile	Gly	Gly	Lys	Lys	Asp	Phe	His	Pro	Glu	Glu	Ile	Ala	Thr	Ala	Leu	
60		545				550					555					560	

6 / 6

Asp Leu Leu Asn Gln His Gly Ser Lys Glu Lys Gln Lys Glu Asp Cys
565 570 575

5 Arg Lys Leu Cys Lys Val Met Thr Thr Gly Gln Ile Ala Ser Glu Glu
580 585 590

Phe Leu Lys Arg Ile Phe Asp Thr Leu Pro Ala Val
595 600

10

 $\langle 210 \rangle$ 3

<211> 45

<212> DNA

<213> Artificial Sequence

15

<220> primer

<223> 5' PCR primer

<400> 3

20

ataagaatgc ggccgccacc atgatccctt ctggcctagt ttatc

45

<210> 4

<211> 34

25

<212> DNA

<213> Artificial Sequence

<220> primer

<223> 3' PCR primer

30

<400> 4

cgggggtaccg aatattcata cagcgggaag tgtg

34

35

INTERNATIONAL SEARCH REPORT

International Application No
PCT/CA 00/01085

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/62 C12N15/85 C12N9/10 C07K16/40
A61K31/711 A61K38/45 A61K39/40 G01N33/53

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 27105 A (GENSET SA ;GRIFFAIS REMY (FR)) 3 June 1999 (1999-06-03) page 127	1-34
X	--- DATABASE SWALL [Online] EBI; 1 May 1999 (1999-05-01) "Lipid A disaccharide synthase (LpxB)" XP002157612 Acc. No. Q9Z6U3 -----	1-34

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 January 2001

Date of mailing of the international search report

24 01. 2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mata Vicente, T.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA 00/01085

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim 27 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

As far as as "in vivo" method is concerned, claim 28 is directed to a diagnostic method practised on the human/animal body and the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 00/01085

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9927105 A	03-06-1999	AU 1170299 A	15-06-1999
		BR 9814878 A	03-10-2000
		EP 1032674 A	06-09-2000
